

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 December 2002 (12.12.2002)

PCT

(10) International Publication Number
WO 02/099099 A2

(51) International Patent Classification⁷: **C12N 15/00**

(21) International Application Number: PCT/EP02/06000

(22) International Filing Date: 31 May 2002 (31.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
01112899.8 1 June 2001 (01.06.2001) EP

(71) Applicant (*for all designated States except US*): **EPIDAU-ROS BIOTECHNOLOGIE AG** [DE/DE]; Am Neuland 1, 82347 Bernried (DE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **PENGER, Anja** [DE/DE]; Kustermannstrasse 22, 82327 Tutzing (DE). **SPRENGER, Reimund** [DE/DE]; Schießstattsiedlung 19, 82362 Weilheim (DE). **BRINKMANN, Ulrich** [DE/DE]; Feichtlstrasse 12, 82362 Weilheim (DE).

(74) Agent: **VOSSIUS & PARTNER**; Siebertstrasse 4, 81675 Munich (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: POLYMORPHISMS IN THE HUMAN GENE FOR CYTOCHROME P450 POLYPEPTIDE 2C8 AND THEIR USE IN DIAGNOSTIC AND THERAPEUTIC APPLICATIONS

(57) Abstract: The present invention relates to a polymorphic CYP2C8-polynucleotide. Moreover, the invention relates to genes or vectors comprising the polynucleotides of the invention and to a host cell genetically engineered with the polynucleotide or gene of the invention. Further, the invention relates to methods for producing molecular variant polypeptides or fragments thereof, methods for producing cells capable of expressing a molecular variant polypeptide and to a polypeptide or fragment thereof encoded by the polynucleotide or the gene of the invention or which is obtainable by the method or from the cells produced by the method of the invention. Furthermore, the invention relates to an antibody which binds specifically the polypeptide of the invention. Moreover, the invention relates to a transgenic non-human animal. The invention also relates to a solid support comprising one or a plurality of the above mentioned polynucleotides, genes, vectors, polypeptides, antibodies or host cells. Furthermore, methods of identifying a polymorphism, identifying and obtaining a prodrug or drug or an inhibitor are also encompassed by the present invention. In addition, the invention relates to methods for producing of a pharmaceutical composition and to methods of diagnosing a disease. Further, the invention relates to a method of detection of the polynucleotide of the invention. Furthermore, comprised by the present invention are a diagnostic and a pharmaceutical composition. Even more, the invention relates to uses of the polynucleotides, genes, vectors, polypeptides or antibodies of the invention. Finally, the invention relates to a diagnostic kit.



WO 02/099099 A2

Polymorphisms in the human gene for cytochrome P450 polypeptide 2C8 and their use in diagnostic and therapeutic applications

The present invention relates to a polymorphic CYP2C8 polynucleotide. Moreover, the invention relates to genes or vectors comprising the polynucleotides of the invention and to a host cell genetically engineered with the polynucleotide or gene of the invention. Further, the invention relates to methods for producing molecular variant polypeptides or fragments thereof, methods for producing cells capable of expressing a molecular variant polypeptide and to a polypeptide or fragment thereof encoded by the polynucleotide or the gene of the invention or which is obtainable by the method or from the cells produced by the method of the invention. Furthermore, the invention relates to an antibody which binds specifically the polypeptide of the invention. Moreover, the invention relates to a transgenic non-human animal. The invention also relates to a solid support comprising one or a plurality of the above mentioned polynucleotides, genes, vectors, polypeptides, antibodies or host cells. Furthermore, methods of identifying a polymorphism, identifying and obtaining a pro-drug or drug or an inhibitor are also encompassed by the present invention. In addition, the invention relates to methods for producing of a pharmaceutical composition and to methods of diagnosing a disease. Further, the invention relates to a method of detection of the polynucleotide of the invention. Furthermore, comprised by the present invention are a diagnostic and a pharmaceutical composition. Even more, the invention relates to uses of the polynucleotides, genes, vectors, polypeptides or antibodies of the invention. Finally, the invention relates to a diagnostic kit.

Cytochrome P450 enzymes are metabolic enzymes differentially expressed in several tissues. Cytochrome P450 2C mRNA was detected in abundance in hepatic tissue, to a lesser extend in extrahepatic tissues, e.g. kidney, adrenal gland, brain, uterus, mammary gland, ovary and duodenum, but neither in testes nor ovary (Klose,

J Biochem Mol Toxicol 13 (1999), 289-95). Of the CYP2C subfamily, clustered on chromosome 10q24.1 (Gray, Genomics 28 (1995), 328-32), CYP2C9 and 2C19 are those which gained major interest due to their prominent role in metabolizing therapeutic drugs. Differential breakdown of their substrates led to the identification of alleles for poor (PM) or extensive metabolizers (EM). Nevertheless, the existence of minor CYP2C8 genes was known and characterized to display about 90% amino acid homology (Goldstein, Pharmacogenetics 4 (1994), 285-99). Only recently, the genomic sequence of CYP2C8, spanning a 31 kb region, was published. Interestingly, the gene is involved in intergenic splicing with CYP2C18 composed of 9 exons (Finta, Genomics 63 (2000), 433-8).

Arachidonic acid is one major endogenous substrate for CYP2C8 and specifically epoxidated to equivalent forms of 11, 12- and 14, 15-epoxides. Concerning xenobiotical substrates CYP2C8 represents the isoform with the narrowest substrate specificity. The anticancer drug taxol (paclitaxel), also well known to be a substrate for MDR-1 (Mechetner, Clin Cancer Res. 4 (1998), 389-398), is known to be the prototype. Several other drugs, e.g. verapamil (Tracy, Br J Clin Pharmacol. 47 (1999), 545-52) and rosiglitazone (Malinowski, Clin Ther. 22 (2000), 1151-68) are preferable substrates for CYP2C8 in comparison to other CYP2Cs or CYP3As. Drugs like benzphetamine, retinoic acid, tolbutamide, benzo(a)pyrene, carbamazepine and R-ibuprofen represent a minor contribution of CYP2C8 (Wrighton, J Clin Invest. 80 (1987), 1017-22; Relling, J Pharmacol Exp Ther. 252 (1990), 442-7; Hamman, Biochem Pharmacol. 54 (1997), 33-41; Kerr, Biochem Pharmacol. 47 (1994), 1969-79; Yun, Cancer Res. 52 (1992), 1868-74; Leo, Arch Biochem Biophys 259 (1987), 241-9). So far, the enzymatic induction has only be observed by phenobarbital and rifampicin (Morel, Eur J Biochem. 191 (1990), 437-44). Thum and Borlak (Thum and Borlak, Br J Pharmacol 130 (2000), 1745-52) found a strong correlation between tissue specific gene expression and enzyme activity. Increased CYP2C8 mRNA expression within the right heart ventricle might explain for the lack of efficacy of cardioselective drugs like verapamil. In a porcine system, Fisslthaler (Fisslthaler, Nature 401(1999), 493-7; Fisslthaler, Semin Perinatol 24 (2000), 15-9; Fisslthaler, Circ Res. 88 (2001), 44-51) could show that CYP2C8 meets all criteria for the coronary endothelium-derived hyperpolarisation factor synthase acting on vascular smooth muscle cells prior to dilation.

Since the mRNA has been published, first single nucleotide polymorphisms (SNPs) in exons 3, 5 and 8 were reported in an abstract (Goldstein, Microsomes and Oxidation, Stresa (2000), Italy): an exchange in position 139 of Arg to Lys (exon 3) could be linked to a SNP in exon 8 (Lys399Arg), occurring primarily in Caucasians, and correlated to poor metabolizing phenotype (PM). Exon 5 displays a mutation (Iso269Phe) that is associated with poor metabolizing enzyme restricted to African-Americans. The regulation of the 2Cs is supposed to be modified by polymorphisms in the untranslated region. Regarding CYP2C8, two previously unidentified transcription regulatory factor sites for C/EBP and HPF-1, but no relevant SNPs were identified by Goldstein and coworkers (Goldstein, Microsomes and Oxidation, Stresa Italy (2000), Italy) in that region.

However, means and methods for reliable and improved diagnosing and treating a variety of diseases and disorders or for predicting and overcoming undesired drug effects or interactions based on dysfunctions or dysregulations of cytochrome 2C8 variants were not available yet but are nevertheless highly desirable. Thus, the technical problem underlying the present invention is to comply with the above specified needs.

The solution to this technical problem is achieved by providing the embodiments characterized in the claims.

Accordingly, the present invention relates to a polynucleotide comprising a polynucleotide selected from the group consisting of:

- (a) a polynucleotide having the nucleic acid sequence of SEQ ID NO: 54, 57, 60, 63, 66, 69, 72, 75, 78, 81, 84, 87, 90, 93, 96, 99, 102, 105, 108, 111, 114, 117, 120, 123, 126, 129, 132, 135, 138, 141, 144, 147, 150, 153, 156, 159, 162, 165, 168, 171, 174, 177, 180, 183, 186, 189, 192, 195, 198, 201, 210, 213, 216, 219, 222, 225, 228, 231, 234, 237, 240, 243, 246, 249, 252, 255, 258, 261, 264, 267, 270, 273, 276, 279, 282, 285, 288, 291, 306, 309, 318, 321, 324, 327, 330, 333, 342, 345, 348, 351, 354, 357, 360, 363, 366, 369, 384, 387, 390, 393, 396 or 399;

- (b) a polynucleotide encoding a polypeptide having the amino acid sequence of SEQ ID NO: 6, 8, 10, 12, 18, 377, 379 or 381;
- (c) a polynucleotide capable of hybridizing to a CYP2C8 gene, wherein said polynucleotide is having at a position corresponding to position 411, 560, 713, 817, 824, 831, 879, 886, 1058, 1627, 1668, 1767, 1887, 1905 or 1952 (GenBank accession No: AF136830.1), at a position corresponding to position 171 or 258 (GenBank accession No: AF136832.1), at a position corresponding to position 122, 150, 182, 334, 339 or 378 (GenBank accession No: AF136833.1), at a position corresponding to position 162, 163, 243 (GenBank accession No: AF136834.2) or at position 583 (GenBank accession No: NM_000770.1), at a position corresponding to position 13 or 180 (GenBank accession No: AF136835.1), at a position corresponding to position 116, 132, 172 or 189 (GenBank accession No: AF136836.1), at a position corresponding to position 42 or 101 (GenBank accession No: AF136837.1), at a position corresponding to position 309 (GenBank accession No: AF136838.1), at a position corresponding to position 1135 (GenBank accession No: NM_000770.1), at a position corresponding to position 232 (GenBank accession No: AF136840.1), at a position corresponding to position 206 (GenBank accession No: AF136842.1), at a position corresponding to position 30, 87, 167, 197, 212, 221, 255 or 271 (GenBank accession No: AF136843.1), at a position corresponding to position 118 (GenBank accession No: AF136844.1), at a position corresponding to position 44 (GenBank accession No: AF136845.1) of the cytochrome 2C8 gene (GenBank accession No: GI: 13787189) a nucleotide substitution, at a position corresponding to position 306 to 307, 1271 to 1273 or 1397 to 1398 of the CYP2C8 gene (GenBank accession No: AF136830.1), at a position corresponding to position 329 of the CYP2C8 gene (GenBank accession No: AF136833.1), at a position corresponding to position 87 of the CYP2C8 gene (GenBank accession No: AF136834.2) a deletion of one or more nucleotides or at a position corresponding to position 1785/1786 of the CYP2C8 gene (GenBank accession No: AF136830.1) or at a position corresponding to position 180/181

- of the CYP2C8 gene (GenBank accession No: AF136833.1) an insertion of one or more nucleotides;
- (d) a polynucleotide capable of hybridizing to a CYP2C8 gene, wherein said polynucleotide is having at a position corresponding to position 411, 817, 824, 831, 879, 1058, 1767 or 1887 of the CYP2C8 gene (GenBank accession No: AF136830.1) an A, at a position corresponding to position 560 or 1668 of the CYP2C8 gene (GenBank accession No: AF136830.1) a G, at a position corresponding to position 713 or 886 of the CYP2C8 gene (GenBank accession No: AF136830.1) a T, at a position corresponding to position 1627, 1905 or 1952 of the CYP2C8 gene (GenBank accession No: AF136830.1) a C, at a position corresponding to position 258 of the CYP2C8 gene (GenBank accession No: AF136832.1) a T, at a position corresponding to position 171 of the CYP2C8 gene (GenBank accession No: AF136832.1) a C, at a position corresponding to position 122, 150 or 334 of the CYP2C8 gene (GenBank accession No: AF136833.1) an A, at a position corresponding to position 182 or 378 of the CYP2C8 gene (GenBank accession No: AF136833.1) a C, at a position corresponding to position 162, 163, 243 [identical to position corresponding to position 583 of the CYP2C8 gene (GenBank accession No: NM_000770.1) of the CYP2C8 gene (GenBank accession No: AF136834.2) an A, at a position corresponding to position 180 of the CYP2C8 gene (GenBank accession No: AF136835.1) an A, at a position corresponding to position 13 of the CYP2C8 gene (GenBank accession No: AF136835.1) a G, at a position corresponding to position 116 or 132 of the CYP2C8 gene (GenBank accession No: AF136836.1) a G, at a position corresponding to position 172 of the CYP2C8 gene (GenBank accession No: AF136836.1) a G, at a position corresponding to position 189 of the CYP2C8 gene (GenBank accession No: AF136836.1) a C, at a position corresponding to position 42 or 101 of the CYP2C8 gene (GenBank accession No: AF136837.1) a G, at a position corresponding to position 1135 of the CYP2C8 gene (GenBank accession No: GI: 13787189) an A, at a position corresponding to position 309 of the CYP2C8 gene (GenBank accession No: AF136838.1) a T, at a position corresponding to position 232 (GenBank accession No: 136840.1) a T, at a position corresponding to position 30 or 212 of the CYP2C8 gene (GenBank

- accession No: AF136843.1) a T, at a position corresponding to position 87 of the CYP2C8 gene (GenBank accession No: AF136843.1) a G, at a position corresponding to position 167 or 197 of the CYP2C8 gene (GenBank accession No: AF136843.1) an A, at a position corresponding to position 221, 255 or 271 of the CYP2C8 gene (GenBank accession No: AF136843.1) a C, at a position corresponding to position 118 of the CYP2C8 gene (GenBank accession No: AF136844.1) an A, at a position corresponding to position 44 of the CYP2C8 gene (GenBank accession No: AF136845.1) a T;
- (e) a polynucleotide encoding a molecular CYP2C8 variant polypeptide or fragment thereof, wherein said polypeptide comprises an amino acid substitution at a position corresponding to any one of position 159, 181, 209, 244, 263, 274, 343 or 365 of the CYP2C8 polypeptide (GI: 13787189); and
- (f) a polynucleotide encoding a molecular CYP2C8 variant polypeptide or fragment thereof, wherein said polypeptide comprises an amino acid substitution of T to P at position corresponding to position 159 (frameshift), V to I at a position corresponding to position 181, N to S at a position corresponding to position 209, I to V at a position corresponding to position 244, F to L at a position corresponding to position 263, E to Stop at a position corresponding to position 274, G to S at a position corresponding to position 365 or S to I at a position corresponding to position 343 of the CYP2C8 polypeptide (GenBank accession No: GI: 13787189).

In the context of the present invention the term “polynucleotides” or the term “polypeptides” refers to different variants of a polynucleotide or polypeptide. Said variants comprise a reference or wild type sequence of the polynucleotides or polypeptides of the invention as well as variants which differ therefrom in structure or composition. Reference or wild type sequences for the polynucleotides are GenBank accession No: NM_000770.1 for mRNA. Reference or wild type sequence for the polynucleotide of the invention is for the 5'UTR: GenBank accession No: AF136830.1; for exon 1: GenBank accession No: AF136831.1; for exon 2: GenBank accession No: AF136832.1 and AF136833.1; for exon 3: GenBank accession No: AF136833.1; for exon 4: GenBank accession No: AF136834.2 and AF136835.1; for exon 5: GenBank accession No: AF136836.1 and AF136837.1; for exon 6: GenBank

accession No: AF136838.1 and AF136839.1; for exon 7: GenBank accession No: AF136840.1 and AF136841.1; for exon 8: GenBank accession No: AF136842.1 and AF136843.1; for exon9/3'UTR: GenBank accession No: AF136844.1 and AF136845.1; partly in combination with NM_000770.1 (mRNA). Reference or wild type sequence for the polypeptide of the CYP2C8 gene is GenBank accession No: GI: 13787189. In the context of the present invention the term "5'UTR" refers to the untranslated region 5' to the ATG start codon including the 5'upstream region encompassing the promoter. The term "3'UTR" refers to the untranslated region 3' to the Stop codon.

The differences in structure or composition usually occur by way of nucleotide or amino acid substitution(s), addition(s) and/or deletion(s). Preferred substitution in accordance with the present invention are a T to G substitution at a position corresponding to position 1668 (GenBank accession No: AF136830.1), a G to A substitution at a position corresponding to position 831 (GenBank accession No: AF136830.1), a G to T substitution at a position corresponding to position 309 (GenBank accession No: AF136838.1) and 232 (GenBank accession No: AF136840.1) of the CYP2C8 gene. Preferred deletions in accordance with the invention are an AT deletion at a position corresponding to position 1397 to 1398 (GenBank accession No: AF136830.1) and a deletion of at least one A at a position corresponding to position 329 (GenBank accession No: AF136833.1) of the CYP2C8 gene.

In accordance with the present invention it has also been found that a deletion of the nucleotide A at a position corresponding to position 329 (GenBank accession No: AF136833.1) of the CYP2C8 gene leads to an altered C-terminus of the protein encoding a CYP2C8 polypeptide wherein said polypeptide comprises an amino acid substitution of T to P at a position corresponding to position 159 of the CYP2C8 polypeptide (GenBank accession No: GI: 13787189). In accordance with the present invention it has also been found that a substitution of a G to a T at a position corresponding to position 309 (GenBank accession No: AF136838.1) of the CYP2C8 gene leads to a polypeptide wherein said polypeptide comprises an amino acid substitution of E to a premature termination (stop) at a position corresponding to position 274 of the CYP2C8 polypeptide (GenBank accession No: GI: 13787189) and a substitution of the nucleotide G to an A at a position corresponding to position 1135

(GenBank accession No: NM_000770.1) of the CYP2C8 gene leads to a polypeptide wherein said polypeptide comprises an amino acid substitution of G to S at a position corresponding to position 365 of the CYP2C8 polypeptide (GenBank accession No: GI: 13787189). This will alter the structure or conformation of the protein and will abolish the activity of the drug metabolizing enzyme.

Preferably, said nucleotide substitution(s), addition(s) or deletion(s) comprised by the present invention result(s) in one or more changes of the corresponding amino acid(s) of the polypeptides of the invention.

The variant polynucleotides and polypeptides also comprise fragments of said polynucleotides or polypeptides of the invention. The polynucleotides and polypeptides as well as the aforementioned fragments thereof of the present invention are characterized as being associated with a CYP2C8 dysfunction or dysregulation comprising, e.g., insufficient and/or altered metabolism. Said dysfunctions or dysregulations referred to in the present invention cause a disease or disorder or a prevalence for said disease or disorder. Preferably, as will be discussed below in detail, said disease is a deficiency in the metabolism of certain drugs which are metabolized by CYP2C8, e.g. Taxol, Verapamil, or any other disease caused by a dysfunction or dysregulation due to a polynucleotide or polypeptides of the invention, also referred to as CYP2C8 gene associated diseases in the following.

The term "hybridizing" as used herein refers to polynucleotides which are capable of hybridizing to the polynucleotides of the invention or parts thereof which are associated with a CYP2C8 dysfunction or dysregulation. Thus, said hybridizing polynucleotides are also associated with said dysfunctions and dysregulations. Preferably, said polynucleotides capable of hybridizing to the polynucleotides of the invention or parts thereof which are associated with CYP2C8 dysfunctions or dysregulations are at least 70%, at least 80%, at least 95% or at least 100% identical to the polynucleotides of the invention or parts thereof which are associated with CYP2C8 dysfunctions or dysregulations. Therefore, said polynucleotides may be useful as probes in Northern or Southern Blot analysis of RNA or DNA preparations, respectively, or can be used as oligonucleotide primers in PCR analysis dependent on their respective size. Also comprised by the invention are hybridizing polynucleotides which are useful for analyzing DNA-Protein interactions via, e.g.,

electrophoretic mobility shift analysis (EMSA). Preferably, said hybridizing polynucleotides comprise at least 10, more preferably at least 15 nucleotides in length while a hybridizing polynucleotide of the present invention to be used as a probe preferably comprises at least 100, more preferably at least 200, or most preferably at least 500 nucleotides in length.

It is well known in the art how to perform hybridization experiments with nucleic acid molecules, i.e. the person skilled in the art knows what hybridization conditions s/he has to use in accordance with the present invention. Such hybridization conditions are referred to in standard text books such as Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory (1989) N.Y. Preferred in accordance with the present inventions are polynucleotides which are capable of hybridizing to the polynucleotides of the invention or parts thereof which are associated with a CYP2C8 dysfunction or dysregulation under stringent hybridization conditions, i.e. which do not cross hybridize to unrelated polynucleotides such as polynucleotides encoding a polypeptide different from the CYP2C8 polypeptides of the invention.

The term "corresponding" as used herein means that a position is not only determined by the number of the preceding nucleotides and amino acids, respectively. The position of a given nucleotide or amino acid in accordance with the present invention which may be deleted, substituted or comprise one or more additional nucleotide(s) may vary due to deletions or additional nucleotides or amino acids elsewhere in the gene or the polypeptide. Thus, under a "corresponding position" in accordance with the present invention it is to be understood that nucleotides or amino acids may differ in the indicated number but may still have similar neighboring nucleotides or amino acids. Said nucleotides or amino acids which may be exchanged, deleted or comprise additional nucleotides or amino acids are also comprised by the term "corresponding position". Said nucleotides or amino acids may for instance together with their neighbors form sequences which may be involved in the regulation of gene expression, stability of the corresponding RNA or RNA editing, as well as encode functional domains or motifs of the protein of the invention.

By, e.g., "position 1271 to 1273" it is meant that said polynucleotide comprises one or more deleted nucleotides which are deleted between positions 1271 and position 1273 of the corresponding wild type version of said polynucleotide. The same applies mutatis mutandis to all other position numbers referred to in the above embodiment which are drafted in the same format.

By, e.g., "position 180/181" it is meant that said polynucleotide comprises one or more additional nucleotide(s) which are inserted between positions 180 and position 181 of the corresponding wild type version of said polynucleotide. The same applies mutatis mutandis to all other position numbers referred to in the above embodiment which are drafted in the same format, i.e. two consecutive position numbers separated by a slash (/).

In accordance with the present invention, the mode and population distribution of genetic variations in the CYP2C8 gene has been analyzed by sequence analysis of relevant regions of the human said gene from many different individuals. It is a well known fact that genomic DNA of individuals, which harbor the individual genetic makeup of all genes, including the CYP2C8 gene, can easily be purified from individual blood samples. These individual DNA samples are then used for the analysis of the sequence composition of the alleles of the CYP2C8 gene that are present in the individual which provided the blood sample. The sequence analysis was carried out by PCR amplification of relevant regions of said genes, subsequent purification of the PCR products, followed by automated DNA sequencing with established methods (e.g. ABI dye terminator cycle sequencing).

One important parameter that had to be considered in the attempt to determine the individual genotypes and identify novel variants of the CYP2C8 gene by direct DNA-sequencing of PCR-products from human blood genomic DNA is the fact that each human harbors (usually, with very few abnormal exceptions) two gene copies of each autosomal gene (diploidy). Because of that, great care had to be taken in the evaluation of the sequences to be able to identify unambiguously not only homozygous sequence variations but also heterozygous variations. The details of the different steps in the identification and characterization of novel polymorphisms in the

CYP2C8 gene (homozygous and heterozygous) are described in the examples below.

Over the past 20 years, genetic heterogeneity has been increasingly recognized as a significant source of variation in drug response. Many scientific communications (Meyer, *Ann. Rev. Pharmacol. Toxicol.* 37 (1997), 269-296 and West, *J. Clin. Pharmacol.* 37 (1997), 635-648) have clearly shown that some drugs work better or may even be highly toxic in some patients than in others and that these variations in patient's responses to drugs can be related to molecular basis. This "pharmacogenomic" concept spots correlations between responses to drugs and genetic profiles of patient's (Marshall, *Nature Biotechnology*, 15 (1997), 954-957; Marshall, *Nature Biotechnology*, 15 (1997), 1249-1252). In this context of population variability with regard to drug therapy, pharmacogenomics has been proposed as a tool useful in the identification and selection of patients which can respond to a particular drug without side effects. This identification/selection can be based upon molecular diagnosis of genetic polymorphisms by genotyping DNA from leukocytes in the blood of patient, for example, and characterization of disease (Bertz, *Clin. Pharmacokinet.* 32 (1997), 210-256; Engel, *J. Chromatogra. B. Biomed. Appl.* 678 (1996), 93-103). For the founders of health care, such as health maintenance organizations in the US and government public health services in many European countries, this pharmacogenomics approach can represent a way of both improving health care and reducing overheads because there is a large cost to unnecessary drugs, ineffective drugs and drugs with side effects.

The mutations in the variant genes of the invention sometime result in amino acid deletion(s), insertion(s) and in particular in substitution(s) either alone or in combination. It is of course also possible to genetically engineer such mutations in wild type genes or other mutant forms. Methods for introducing such modifications in the DNA sequence of said genes are well known to the person skilled in the art; see, e.g., Sambrook, *Molecular Cloning A Laboratory Manual*, Cold Spring Harbor Laboratory (1989) N.Y.

For the investigation of the nature of the alterations in the amino acid sequence of

the polypeptides of the invention may be used such as BRASMOL that are obtainable from the Internet. Furthermore, folding simulations and computer redesign of structural motifs can be performed using other appropriate computer programs (Olszewski, *Proteins* 25 (1996), 286-299; Hoffman, *Comput. Appl. Biosci.* 11 (1995), 675-679). Computers can be used for the conformational and energetic analysis of detailed protein models (Monge, *J. Mol. Biol.* 247 (1995), 995-1012; Renouf, *Adv. Exp. Med. Biol.* 376 (1995), 37-45). These analysis can be used for the identification of the influence of a particular mutation on metabolizing, binding and/or transport of drugs.

Usually, said amino acid deletion, addition or substitution in the amino acid sequence of the protein encoded by the polynucleotide of the invention is due to one or more nucleotide substitution, insertion or deletion, or any combinations thereof. Preferably said nucleotide substitution, insertion or deletion may result in an amino acid substitution of F to L at position corresponding to position 263 of the CYP2C8 polypeptide (GenBank accession No: GI: 13787189). The polypeptides encoded by the polynucleotides of the invention have altered biological or immunological properties due to the mutations referred to in accordance with the present invention. Examples for said altered properties are stability of the polypeptides which may be effected or an altered substrate specificity or even a complete loss of the capability of metabolizing certain drugs.

The mutations in the CYP2C8 gene detected in accordance with the present invention are listed in Table 2. The methods of the mutation analysis followed standard protocols and are described in detail in the Examples. In general such methods are to be used in accordance with the present invention for evaluating the phenotypic spectrum as well as the overlapping clinical characteristics of diseases or conditions related to dysfunctions or dysregulations and diseases related to the poor or extensive metabolism (PM or EM) certain drugs. Advantageously, the characterization of said mutants may form the basis of the development of a diagnostic assay that is able to predict a patients efficacy to metabolize a drug for instance in anticancer treatment (taxol) or cardiovascular deficiencies (verapamil). Said methods encompass for example haplotype analysis, single-strand

conformation polymorphism analysis (SSCA), PCR and direct sequencing. On the basis of thorough clinical characterization of many patients the phenotypes can then be correlated to these mutations.

Also comprised by the polynucleotides referred to in the present invention are polynucleotides which comprise at least two of the polynucleotides specified herein above, i.e. polynucleotides having a nucleotide sequence which contains all four mutations comprised by the above polynucleotides or listed in Table 2 below (haplotype: positions 831, 1397 to 1398 of GenBank accession No: AF136830.1, position 270 of GenBank accession No: AF136833.1, and position 206 of (GenBank accession No: AF136842.1). In accordance with the present invention it is also preferred to detect only one of the above mentioned polymorphisms of the haplotype, said one polymorphism being indicative for the presence of the other polymorphisms of the haplotype. Thus, in order to detect the presence of the above mentioned haplotype it is sufficient to determine the presence of any one of the polymorphisms comprised by said haplotype. Moreover, the polynucleotides referred to above allow the study of synergistic effects of said mutations in the CYP2C8 gene and/or a polypeptide encoded by said polynucleotide on the pharmacological profile of drugs in patients who bear such mutant forms of the gene or similar mutant forms that can be mimicked by the above described proteins. It is expected that the analysis of said synergistic effects provides deeper insights into the onset of CYP2C8 dysfunctions or dysregulations or diseases related to altered drug transport as described supra. From said deeper insight the development of diagnostic and pharmaceutical compositions related to CYP2C8 dysfunctions or dysregulations or diseases related to impaired drug metabolism will greatly benefit.

As is evident to the person skilled in the art, the genetic knowledge deduced from the present invention can now be used to exactly and reliably characterize the genotype of a patient. Advantageously, diseases or a prevalence for a disease which are associated with CYP2C8 dysfunction or dysregulation, e.g. diseases associated with arachidonic acid metabolism referred to herein can be predicted and preventive or therapeutical measures can be applied accordingly. Moreover in accordance with the foregoing, in cases where a given drug takes an unusual effect, a suitable individual

therapy can be designed based on the knowledge of the individual genetic makeup of a subject with respect to the polynucleotides of the invention and improved therapeutics can be developed as will be further discussed below.

In general, the CYP2C8 "status", defined by the expression level and activity of the CYP2C8 protein, can be variable in normal tissue, due to genetic variations/polymorphisms. The identification of polymorphisms associated with altered CYP2C8 expression and/or activity is important for the prediction of drug metabolism and subsequently for the prediction of therapy outcome, including side effects of medications. Therefore, analysis of CYP2C8 variations indicative of CYP2C8 function, is a valuable tool for therapy with drugs, which are substrates of CYP2C8 and has, thanks to the present invention, now become possible.

In line with the foregoing, preferably, the polynucleotide of the present invention is associated with an incompatibility or a disease related to arachidonic acid metabolism, cancer or cardiovascular diseases.

The term "cancer" used herein is very well known and characterized in the art. Several variants of cancer exist and are comprised by said term as meant in accordance with the invention. For a detailed list of symptoms which are indicative for cancer it is referred to text book knowledge, e.g. Pschyrembel. The term "cardiovascular disease" as used herein refers to those diseases known in the art and described in detail in standard text books, such as Pschyrembel or Stadman. Examples for cardiovascular diseases are hypertension or atherosclerosis. The inefficacy or complete loss to epoxidate arachidonic acid is referred to as disease of the arachidonic acid metabolism.

In a further embodiment the present invention relates to a polynucleotide which is DNA or RNA.

The polynucleotide of the invention may be, e.g., DNA, cDNA, genomic DNA, RNA or synthetically produced DNA or RNA or a recombinantly produced chimeric nucleic acid molecule comprising any of those polynucleotides either alone or in combination. Preferably said polynucleotide is part of a vector, particularly plasmids, cosmids, viruses and bacteriophages used conventionally in genetic engineering that

comprise a polynucleotide of the invention. Such vectors may comprise further genes such as marker genes which allow for the selection of said vector in a suitable host cell and under suitable conditions.

The invention furthermore relates to a gene comprising the polynucleotide of the invention.

It is well known in the art that genes comprise structural elements which encode an amino acid sequence as well as regulatory elements which are involved in the regulation of the expression of said genes. Structural elements are represented by exons which may either encode an amino acid sequence or which may encode for RNA which is not encoding an amino acid sequence but is nevertheless involved in RNA function, e.g. by regulating the stability of the RNA or the nuclear export of the RNA.

Regulatory elements of a gene may comprise promoter elements or enhancer elements both of which could be involved in transcriptional control of gene expression. It is very well known in the art that a promoter is to be found upstream of the structural elements of a gene. Regulatory elements such as enhancer elements, however, can be found distributed over the entire locus of a gene. Said elements could be reside, e.g., in introns, regions of genomic DNA which separate the exons of a gene. Promoter or enhancer elements correspond to polynucleotide fragments which are capable of attracting or binding polypeptides involved in the regulation of the gene comprising said promoter or enhancer elements. For example, polypeptides involved in regulation of said gene comprise the so called transcription factors.

Said introns may comprise further regulatory elements which are required for proper gene expression. Introns are usually transcribed together with the exons of a gene resulting in a nascent RNA transcript which contains both, exon and intron sequences. The intron encoded RNA sequences are usually removed by a process known as RNA splicing. However, said process also requires regulatory sequences present on a RNA transcript said regulatory sequences may be encoded by the introns.

In addition, besides their function in transcriptional control and control of proper RNA processing and/or stability, regulatory elements of a gene could be also involved in the control of genetic stability of a gene locus. Said elements control, e.g.,

recombination events or serve to maintain a certain structure of the DNA or the arrangement of DNA in a chromosome.

Therefore, single nucleotide polymorphisms can occur in exons of a gene which encode an amino acid sequence as discussed supra as well as in regulatory regions which are involved in the above discussed process. The analysis of the nucleotide sequence of a gene locus in its entirety including, e.g., introns is in light of the above desirable. The polymorphisms comprised by the polynucleotides of the present invention can influence the expression level of CYP2C8 protein via mechanisms involving enhanced or reduced transcription of the CYP2C8 gene, stabilization of the gene's RNA transcripts and alteration of the processing of the primary RNA transcripts.

Therefore, in a furthermore preferred embodiment of the gene of the invention a nucleotide deletion, addition and/or substitution results in altered expression of the variant gene compared to the corresponding wild type gene.

In another embodiment the present invention relates to a vector comprising the polynucleotide of the invention or the gene of the invention.

Said vector may be, for example, a phage, plasmid, viral or retroviral vector. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host/cells.

The polynucleotides or genes of the invention may be joined to a vector containing selectable markers for propagation in a host. Generally, a plasmid vector is introduced in a precipitate such as a calcium phosphate precipitate, or in a complex with a charged lipid or in carbon-based clusters. Should the vector be a virus, it may be packaged in vitro using an appropriate packaging cell line prior to application to host cells.

In a more preferred embodiment of the vector of the invention the polynucleotide is operatively linked to expression control sequences allowing expression in prokaryotic or eukaryotic cells or isolated fractions thereof.

Expression of said polynucleotide comprises transcription of the polynucleotide, preferably into a translatable mRNA. Regulatory elements ensuring expression in

eukaryotic cells, preferably mammalian cells, are well known to those skilled in the art. They usually comprise regulatory sequences ensuring initiation of transcription and optionally poly-A signals ensuring termination of transcription and stabilization of the transcript. Additional regulatory elements may include transcriptional as well as translational enhancers. Possible regulatory elements permitting expression in prokaryotic host cells comprise, e.g., the *lac*, *trp* or *tac* promoter in *E. coli*, and examples for regulatory elements permitting expression in eukaryotic host cells are the *AOX1* or *GAL1* promoter in yeast or the CMV-, SV40-, RSV-promoter (Rous sarcoma virus), CMV-enhancer, SV40-enhancer or a globin intron in mammalian and other animal cells. Beside elements which are responsible for the initiation of transcription such regulatory elements may also comprise transcription termination signals, such as the SV40-poly-A site or the tk-poly-A site, downstream of the polynucleotide. In this context, suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pCDM8, pRc/CMV, pcDNA1, pcDNA3 (In-vitro gene), pSPORT1 (GIBCO BRL). Preferably, said vector is an expression vector and/or a gene transfer or targeting vector. Expression vectors derived from viruses such as retroviruses, vaccinia virus, adeno-associated virus, herpes viruses, or bovine papilloma virus, may be used for delivery of the polynucleotides or vector of the invention into targeted cell population. Methods which are well known to those skilled in the art can be used to construct recombinant viral vectors; see, for example, the techniques described in Sambrook, Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory (1989) N.Y. and Ausubel, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y. (1994). Alternatively, the polynucleotides and vectors of the invention can be reconstituted into liposomes for delivery to target cells.

The term "isolated fractions thereof" refers to fractions of eukaryotic or prokaryotic cells or tissues which are capable of transcribing or transcribing and translating RNA from the vector of the invention. Said fractions comprise proteins which are required for transcription of RNA or transcription of RNA and translation of said RNA into a polypeptide. Said isolated fractions may be, e.g., nuclear and cytoplasmic fractions of eukaryotic cells such as of reticulocytes.

The present invention furthermore relates to a host cell genetically engineered with the polynucleotide of the invention, the gene of the invention or the vector of the invention.

Said host cell may be a prokaryotic or eukaryotic cell; see supra. The polynucleotide or vector of the invention which is present in the host cell may either be integrated into the genome of the host cell or it may be maintained extrachromosomally. In this respect, it is also to be understood that the recombinant DNA molecule of the invention can be used for "gene targeting" and/or "gene replacement", for restoring a mutant gene or for creating a mutant gene via homologous recombination; see for example Mouellic, Proc. Natl. Acad. Sci. USA, 87 (1990), 4712-4716; Joyner, Gene Targeting, A Practical Approach, Oxford University Press.

The host cell can be any prokaryotic or eukaryotic cell, such as a bacterial, insect, fungal, plant, animal, mammalian or, preferably, human cell. Preferred fungal cells are, for example, those of the genus *Saccharomyces*, in particular those of the species *S. cerevisiae*. The term "prokaryotic" is meant to include all bacteria which can be transformed or transfected with a polynucleotide for the expression of a variant polypeptide of the invention. Prokaryotic hosts may include gram negative as well as gram positive bacteria such as, for example, *E. coli*, *S. typhimurium*, *Serratia marcescens* and *Bacillus subtilis*. A polynucleotide coding for a mutant form of variant polypeptides of the invention can be used to transform or transfect the host using any of the techniques commonly known to those of ordinary skill in the art. Methods for preparing fused, operably linked genes and expressing them in bacteria or animal cells are well-known in the art (Sambrook, supra). The genetic constructs and methods described therein can be utilized for expression of variant polypeptides of the invention in, e.g., prokaryotic hosts. In general, expression vectors containing promoter sequences which facilitate the efficient transcription of the inserted polynucleotide are used in connection with the host. The expression vector typically contains an origin of replication, a promoter, and a terminator, as well as specific genes which are capable of providing phenotypic selection of the transformed cells. The transformed prokaryotic hosts can be grown in fermentors and cultured according to techniques known in the art to achieve optimal cell growth. The proteins of the invention can then be isolated from the grown medium, cellular lysates, or cellular membrane fractions. The isolation and purification of the microbially or

otherwise expressed polypeptides of the invention may be by any conventional means such as, for example, preparative chromatographic separations and immunological separations such as those involving the use of monoclonal or polyclonal antibodies.

Thus, in a further embodiment the invention relates to a method for producing a molecular variant CYP2C8 polypeptide or fragment thereof comprising culturing the above described host cell; and recovering said protein or fragment from the culture.

In another embodiment the present invention relates to a method for producing cells capable of expressing a molecular variant CYP2C8 polypeptide comprising genetically engineering cells with the polynucleotide of the invention, the gene of the invention or the vector of the invention.

The cells obtainable by the method of the invention can be used, for example, to test drugs according to the methods described in D. L. Spector, R. D. Goldman, L. A. Leinwand, Cells, a Lab manual, CSH Press 1998. Furthermore, the cells can be used to study known drugs and unknown derivatives thereof for their ability to complement the deficiency caused by mutations in the CYP2C8 gene. For these embodiments the host cells preferably lack a wild type allele, preferably both alleles of the CYP2C8 gene and/or have at least one mutated from thereof. Ideally, the gene comprising an allele as comprised by the polynucleotides of the invention could be introduced into the wild type locus by homologous replacement. Alternatively, strong overexpression of a mutated allele over the normal allele and comparison with a recombinant cell line overexpressing the normal allele at a similar level may be used as a screening and analysis system. The cells obtainable by the above-described method may also be used for the screening methods referred to herein below.

Furthermore, the invention relates to a polypeptide or fragment thereof encoded by the polynucleotide of the invention, the gene of the invention or obtainable by the method described above or from cells produced by the method described above.

In this context it is also understood that the variant polypeptide of the invention can be further modified by conventional methods known in the art. By providing said variant proteins according to the present invention it is also possible to determine the

portions relevant for their biological activity or inhibition of the same. The terms "polypeptide" and "protein" as used herein are exchangeable. Moreover, what is comprised by said terms is standard textbook knowledge.

The present invention furthermore relates to an antibody which binds specifically to the polypeptide of the invention.

Advantageously, the antibody specifically recognizes or binds an epitope containing one or more amino acid substitution(s) as defined above. Antibodies against the variant polypeptides of the invention can be prepared by well known methods using a purified protein according to the invention or a (synthetic) fragment derived therefrom as an antigen. Monoclonal antibodies can be prepared, for example, by the techniques as originally described in Köhler and Milstein, *Nature* 256 (1975), 495, and Galfré, *Meth. Enzymol.* 73 (1981), 3, which comprise the fusion of mouse myeloma cells to spleen cells derived from immunized mammals. In a preferred embodiment of the invention, said antibody is a monoclonal antibody, a polyclonal antibody, a single chain antibody, human or humanized antibody, primatized, chimerized or fragment thereof that specifically binds said peptide or polypeptide also including bispecific antibody, synthetic antibody, antibody fragment, such as Fab, Fv or scFv fragments etc., or a chemically modified derivative of any of these. Furthermore, antibodies or fragments thereof to the aforementioned polypeptides can be obtained by using methods which are described, e.g., in Harlow and Lane "Antibodies, A Laboratory Manual", CSH Press, Cold Spring Harbor, 1988. These antibodies can be used, for example, for the immunoprecipitation and immunolocalization of the variant polypeptides of the invention as well as for the monitoring of the presence of said variant polypeptides, for example, in recombinant organisms, and for the identification of compounds interacting with the proteins according to the invention. For example, surface plasmon resonance as employed in the BIAcore system can be used to increase the efficiency of phage antibodies which bind to an epitope of the protein of the invention (Schier, *Human Antibodies Hybridomas* 7 (1996), 97-105; Malmberg, *J. Immunol. Methods* 183 (1995), 7-13).

In a preferred embodiment the antibody of the present invention specifically recognizes an epitope containing one or more amino acid substitution(s) resulting from a nucleotide exchange as defined supra.

Antibodies which specifically recognize modified amino acids such as phospho-Tyrosine residues are well known in the art. Similarly, in accordance with the present invention antibodies which specifically recognize even a single amino acid exchange in an epitope may be generated by the well known methods described supra.

In light of the foregoing, in a more preferred embodiment the antibody of the present invention is monoclonal or polyclonal.

The invention also relates to a transgenic non-human animal comprising at least one polynucleotide of the invention, the gene of the invention or the vector of the invention as described supra.

The present invention also encompasses a method for the production of a transgenic non-human animal comprising introduction of a polynucleotide or vector of the invention into a germ cell, an embryonic cell, stem cell or an egg or a cell derived therefrom. The non-human animal can be used in accordance with the method of the invention described below and may be a non-transgenic healthy animal, or may have a disease or disorder, preferably a disease caused by at least one mutation in the gene of the invention. Such transgenic animals are well suited for, e.g., pharmacological studies of drugs in connection with variant forms of the above described variant polypeptides since these polypeptides or at least their functional domains are conserved between species in higher eukaryotes, particularly in mammals. Production of transgenic embryos and screening of those can be performed, e.g., as described by A. L. Joyner Ed., Gene Targeting, A Practical Approach (1993), Oxford University Press. The DNA of the embryos can be analyzed using, e.g., Southern blots with an appropriate probe or based on PCR techniques.

A transgenic non-human animal in accordance with the invention may be a transgenic mouse, rat, hamster, dog, monkey, rabbit, pig, frog, nematode such as *Caenorhabditis elegans*, fruitfly such as *Drosophila melanogaster* or fish such as torpedo fish or zebrafish comprising a polynucleotide or vector of the invention or obtained by the method described above, preferably wherein said polynucleotide or

vector is stably integrated into the genome of said non-human animal, preferably such that the presence of said polynucleotide or vector leads to the expression of the variant polypeptide of the invention. It may comprise one or several copies of the same or different polynucleotides or genes of the invention. This animal has numerous utilities, including as a research model for cardiovascular research and therefore, presents a novel and valuable animal in the development of therapies, treatment, etc. for diseases caused by cardiovascular diseases. Accordingly, in this instance, the mammal is preferably a laboratory animal such as a mouse or rat.

Thus, in a preferred embodiment the transgenic non-human animal of the invention is a mouse, a rat or a zebrafish.

Numerous reports revealed that said animals are particularly well suited as model organisms for the investigation of the drug metabolism and its deficiencies or cancer. Advantageously, transgenic animals can be easily created using said model organisms, due to the availability of various suitable techniques well known in the art.

The invention also relates to a solid support comprising one or a plurality of the polynucleotide, the gene, the vector, the polypeptide, the antibody or the host cell of the invention in immobilized form.

The term "solid support" as used herein refers to a flexible or non-flexible support that is suitable for carrying said immobilized targets. Said solid support may be homogenous or inhomogeneous. For example, said solid support may consist of different materials having the same or different properties with respect to flexibility and immobilization, for instance, or said solid support may consist of one material exhibiting a plurality of properties also comprising flexibility and immobilization properties. Said solid support may comprise glass-, polypropylene- or silicon-chips, membranes oligonucleotide-conjugated beads or bead arrays.

The term "immobilized" means that the molecular species of interest is fixed to a solid support, preferably covalently linked thereto. This covalent linkage can be achieved by different means depending on the molecular nature of the molecular species. Moreover, the molecular species may be also fixed on the solid support by electrostatic forces, hydrophobic or hydrophilic interactions or Van-der-Waals forces. The above described physico-chemical interactions typically occur in interactions

between molecules. For example, biotinylated polypeptides may be fixed on a avidin-coated solid support due to interactions of the above described types. Further, polypeptides such as antibodies, may be fixed on an antibody coated solid support. Moreover, the immobilization is dependent on the chemical properties of the solid support. For example, the nucleic acid molecules can be immobilized on a membrane by standard techniques such as UV-crosslinking or heat.

In a preferred embodiment of the invention said solid support is a membrane, a glass- or polypropylene- or silicon-chip, are oligonucleotide-conjugated beads or a bead array, which is assembled on an optical filter substrate.

Moreover, the present invention relates to an in vitro method for identifying a polymorphism said method comprising the steps of:

- (a) isolating a polynucleotide or the gene of the invention from a plurality of subgroups of individuals, wherein one subgroup has no prevalence for a CYP2C8 associated disease and at least one or more further subgroup(s) do have prevalence for a CYP2C8 associated disease; and
- (b) identifying a polymorphism by comparing the nucleic acid sequence of said polynucleotide or said gene of said one subgroup having no prevalence for a CYP2C8 associated disease with said at least one or more further subgroup(s) having a prevalence for a CYP2C8 associated disease.

The term "prevalence" as used herein means that individuals are be susceptible for one or more disease(s) which are associated with CYP2C8 dysfunction or dysregulation or could already have one or more of said disease(s). Thereby, one CYP2C8 associated disease can be used to determine the susceptibility for another CYP2C8 associated disease. Moreover, symptoms which are indicative for a prevalence for developing of a disease are very well known in the art and have been sufficiently described in standard textbooks such as Pschyrembel.

Advantageously, polymorphisms according to the present invention which are associated with CYP2C8 dysfunction or dysregulation or one or more disease(s) based thereon should be enriched in subgroups of individuals which have a prevalence for said diseases versus subgroups which have no prevalence for said

diseases. Thus, the above described method allows the rapid and reliable detection of polymorphism which are indicative for one or more CYP2C8 associated disease(s) or a susceptibility therefor. Advantageously, due to the phenotypic preselection a large number of individuals having no prevalence might be screened for polymorphisms in general. Thereby, a reference sequences comprising polymorphisms which do not correlate to one or more CYP2C8 associated disease(s) can be obtained. Based on said reference sequences it is possible to efficiently and reliably determine the relevant polymorphisms.

In a further embodiment the present invention relates to a method for identifying and obtaining a pro-drug or a drug capable of modulating the activity of a molecular variant of a CYP2C8 polypeptide comprising the steps of:

- (a) contacting the polypeptide, the solid support of the invention, a cell expressing a molecular variant gene comprising a polynucleotide of the invention, the gene or the vector of the invention in the presence of components capable of providing a detectable signal in response to drug activity with a compound to be screened for pro-drug or drug activity; and
- (b) detecting the presence or absence of a signal or increase or decrease of a signal generated from the pro-drug or the drug activity, wherein the absence, presence, increase or decrease of the signal is indicative for a putative pro-drug or drug.

The term "compound" in a method of the invention includes a single substance or a plurality of substances which may or may not be identical.

Said compound(s) may be chemically synthesized or produced via microbial fermentation but can also be comprised in, for example, samples, e.g., cell extracts from, e.g., plants, animals or microorganisms. Furthermore, said compounds may be known in the art but hitherto not known to be useful as an inhibitor, respectively. The plurality of compounds may be, e.g., added to the culture medium or injected into a cell or non-human animal of the invention.

If a sample containing (a) compound(s) is identified in the method of the invention, then it is either possible to isolate the compound from the original sample identified as containing the compound, in question or one can further subdivide the original

sample, for example, if it consists of a plurality of different compounds, so as to reduce the number of different substances per sample and repeat the method with the subdivisions of the original sample. It can then be determined whether said sample or compound displays the desired properties, for example, by the methods described herein or in the literature (Spector et al., Cells manual; see supra). Depending on the complexity of the samples, the steps described above can be performed several times, preferably until the sample identified according to the method of the invention only comprises a limited number of or only one substance(s). Preferably said sample comprises substances of similar chemical and/or physical properties, and most preferably said substances are identical. The methods of the present invention can be easily performed and designed by the person skilled in the art, for example in accordance with other cell based assays described in the prior art or by using and modifying the methods as described herein. Furthermore, the person skilled in the art will readily recognize which further compounds may be used in order to perform the methods of the invention, for example, enzymes, if necessary, that convert a certain compound into a precursor. Such adaptation of the method of the invention is well within the skill of the person skilled in the art and can be performed without undue experimentation.

Compounds which can be used in accordance with the present invention include peptides, proteins, nucleic acids, antibodies, small organic compounds, ligands, peptidomimetics, PNAs and the like. Said compounds may act as agonists or antagonists of the invention. Said compounds can also be functional derivatives or analogues of known drugs. Methods for the preparation of chemical derivatives and analogues are well known to those skilled in the art and are described in, for example, Beilstein, Handbook of Organic Chemistry, Springer edition New York Inc., 175 Fifth Avenue, New York, N.Y. 10010 U.S.A. and Organic Synthesis, Wiley, New York, USA. Furthermore, said derivatives and analogues can be tested for their effects according to methods known in the art or as described. Furthermore, peptide mimetics and/or computer aided design of appropriate drug derivatives and analogues can be used, for example, according to the methods described below. Such analogs comprise molecules that may have the basis structure of known CYP2C8 substrates, inhibitors and/or modulators.

Appropriate computer programs can be used for the identification of interactive sites of a putative inhibitor and the polypeptides of the invention by computer assisted searches for complementary structural motifs (Fassina, Immunomethods 5 (1994), 114-120). Further appropriate computer systems for the computer aided design of protein and peptides are described in the prior art, for example, in Berry, Biochem. Soc. Trans. 22 (1994), 1033-1036; Wodak, Ann. N. Y. Acad. Sci. 501 (1987), 1-13; Pabo, Biochemistry 25 (1986), 5987-5991. The results obtained from the above-described computer analysis can be used in combination with the method of the invention for, e.g., optimizing known inhibitors, analogs, antagonists or agonists. Appropriate peptidomimetics and other inhibitors can also be identified by the synthesis of peptidomimetic combinatorial libraries through successive chemical modification and testing the resulting compounds, e.g., according to the methods described herein. Methods for the generation and use of peptidomimetic combinatorial libraries are described in the prior art, for example in Ostresh, Methods in Enzymology 267 (1996), 220-234 and Dorner, Bioorg. Med. Chem. 4 (1996), 709-715. Furthermore, the three-dimensional and/or crystallographic structure of said compounds and the polypeptides of the invention can be used for the design of peptidomimetic drugs (Rose, Biochemistry 35 (1996), 12933-12944; Rutenber, Bioorg. Med. Chem. 4 (1996), 1545-1558). It is very well known how to obtain said compounds, e.g. by chemical or biochemical standard techniques. Thus, also comprised by the method of the invention are means of making or producing said compounds. In summary, the present invention provides methods for identifying and obtaining compounds which can be used in specific doses for the treatment of specific forms of CYP2C8 associated diseases.

The above definitions apply mutatis mutandis to all of the methods described in the following.

In a further embodiment the present invention relates to a method for identifying and obtaining an inhibitor of the activity of a molecular variant of a CYP2C8 polypeptide comprising the steps of:

- (a) contacting the protein, the solid support of the invention or a cell expressing a molecular variant gene comprising a polynucleotide or the gene or the vector

of the invention in the presence of components capable of providing a detectable signal in response to drug activity with a compound to be screened for inhibiting activity; and

- (b) detecting the presence or absence of a signal or increase or decrease of a signal generated from the inhibiting activity, wherein the absence or decrease of the signal is indicative for a putative inhibitor.

In a preferred embodiment of the method of the invention said cell is a cell, obtained by the method of the invention or can be obtained from the transgenic non-human animal as described supra.

In a still further embodiment the present invention relates to a method of identifying and obtaining a pro-drug or drug capable of modulating the activity of a molecular variant of a CYP2C8 polypeptide comprising the steps of:

- (a) contacting the host cell, the cell obtained by the method of the invention, the polypeptide or the solid support of the invention with the first molecule known to be bound by a CYP2C8 polypeptide to form a first complex of said polypeptide and said first molecule;
- (b) contacting said first complex with a compound to be screened, and
- (c) measuring whether said compound displaces said first molecule from said first complex.

Advantageously, in said method said measuring step comprises measuring the formation of a second complex of said protein and said inhibitor candidate. Preferably, said measuring step comprises measuring the amount of said first molecule that is not bound to said protein.

In a particularly preferred embodiment of the above-described method of said first molecule is a agonist or antagonist or a substrate and/or a inhibitor and/or a modulator of the polypeptide of the invention, e.g., with a radioactive or fluorescent label.

In a still another embodiment the present invention relates to a method of identifying and obtaining an inhibitor capable of modulating the activity of a molecular variant of a CYP2C8 polypeptide comprising the steps of:

- (a) contacting the host cell or the cell obtained by the method of the invention, the protein or the solid support of the invention with the first molecule known to be bound by the CYP2C8 polypeptide to form a first complex of said protein and said first molecule;
- (b) contacting said first complex with a compound to be screened, and
- (c) measuring whether said compound displaces said first molecule from said first complex.

In a preferred embodiment of the method of the invention said measuring step comprises measuring the formation of a second complex of said protein and said compound.

In another preferred embodiment of the method of the invention said measuring step comprises measuring the amount of said first molecule that is not bound to said protein.

In a more preferred embodiment of the method of the invention said first molecule is labeled.

The invention furthermore relates to a method for the production of a pharmaceutical composition comprising the steps of the method as described supra; and the further step of formulating the compound identified and obtained or a derivative thereof in a pharmaceutically acceptable form.

The therapeutically useful compounds identified according to the methods of the invention can be formulated and administered to a patient as discussed above. For uses and therapeutic doses determined to be appropriate by one skilled in the art and for definitions of the term "pharmaceutical composition" see *infra*.

Furthermore, the present invention encompasses a method for the preparation of a pharmaceutical composition comprising the steps of the above-described methods; and formulating a drug or pro-drug in the form suitable for therapeutic application and

preventing or ameliorating the disorder of the subject diagnosed in the method of the invention.

Drugs or pro-drugs after their *in vivo* administration are metabolized in order to be eliminated either by excretion or by metabolism to one or more active or inactive metabolites (Meyer, J. Pharmacokinet. Biopharm. 24 (1996), 449-459). Thus, rather than using the actual compound or inhibitor identified and obtained in accordance with the methods of the present invention a corresponding formulation as a pro-drug can be used which is converted into its active in the patient. Precautionary measures that may be taken for the application of pro-drugs and drugs are described in the literature; see, for review, Ozama, J. Toxicol. Sci. 21 (1996), 323-329).

In a preferred embodiment of the method of the present invention said drug or prodrug is a derivative of a medicament as defined hereinafter.

The present invention also relates to a method of diagnosing a disorder related to the presence of a molecular variant of the CYP2C8 gene or susceptibility to such a disorder comprising determining the presence of a polynucleotide or the gene of the invention in a sample from a subject.

In accordance with this embodiment of the present invention, the method of testing the status of a disorder or susceptibility to such a disorder can be effected by using a polynucleotide gene or nucleic acid of the invention, e.g., in the form of a Southern or Northern blot or *in situ* analysis. Said nucleic acid sequence may hybridize to a coding region of either of the genes or to a non-coding region, e.g. intron. In the case that a complementary sequence is employed in the method of the invention, said nucleic acid molecule can again be used in Northern blots. Additionally, said testing can be done in conjunction with an actual blocking, e.g., of the transcription of the gene and thus is expected to have therapeutic relevance. Furthermore, a primer or oligonucleotide can also be used for hybridizing to one of the above mentioned CYP2C8 gene or corresponding mRNAs. The nucleic acids used for hybridization can, of course, be conveniently labeled by incorporating or attaching, e.g., a radioactive or other marker. Such markers are well known in the art. The labeling of said nucleic acid molecules can be effected by conventional methods.

Additionally, the presence or expression of variant CYP2C8 gene can be monitored by using a primer pair that specifically hybridizes to either of the corresponding

nucleic acid sequences and by carrying out a PCR reaction according to standard procedures. Specific hybridization of the above mentioned probes or primers preferably occurs at stringent hybridization conditions. The term "stringent hybridization conditions" is well known in the art; see, for example, Sambrook et al., "Molecular Cloning, A Laboratory Manual" second ed., CSH Press, Cold Spring Harbor, 1989; "Nucleic Acid Hybridisation, A Practical Approach", Hames and Higgins eds., IRL Press, Oxford, 1985. Furthermore, the mRNA, cRNA, cDNA or genomic DNA obtained from the subject may be sequenced to identify mutations which may be characteristic fingerprints of mutations in the polynucleotide or the gene of the invention. The present invention further comprises methods wherein such a fingerprint may be generated by RFLPs of DNA or RNA obtained from the subject, optionally the DNA or RNA may be amplified prior to analysis, the methods of which are well known in the art. RNA fingerprints may be performed by, for example, digesting an RNA sample obtained from the subject with a suitable RNA-Enzyme, for example RNase T₁, RNase T₂ or the like or a ribozyme and, for example, electrophoretically separating and detecting the RNA fragments as described above. Further modifications of the above-mentioned embodiment of the invention can be easily devised by the person skilled in the art, without any undue experimentation from this disclosure; see, e.g., the examples. An additional embodiment of the present invention relates to a method wherein said determination is effected by employing an antibody of the invention or fragment thereof. The antibody used in the method of the invention may be labeled with detectable tags such as a histidine flags or a biotin molecule.

The invention relates to a method of diagnosing a disorder related to the presence of a molecular variant of a CYP2C8 gene or susceptibility to such a disorder comprising determining the presence of a polypeptide or the antibody of the invention in a sample from a subject.

In a preferred embodiment of the above described method said disorder is a cancer or cardiovascular disease.

In a preferred embodiment of the present invention, the above described method is comprising PCR, ligase chain reaction, restriction digestion, direct sequencing, nucleic acid amplification techniques, hybridization techniques or immunoassays.

Said techniques are very well known in the art.

Moreover, the invention relates to a method of detection of the polynucleotide or the gene of the invention in a sample comprising the steps of

- (a) contacting the solid support described supra with the sample under conditions allowing interaction of the polynucleotide or the gene of the invention with the immobilized targets on a solid support and;
- (b) determining the binding of said polynucleotide or said gene to said immobilized targets on a solid support.

The invention also relates to an in vitro method for diagnosing a disease comprising the steps of the method described supra, wherein binding of said polynucleotide or gene to said immobilized targets on said solid support is indicative for the presence or the absence of said disease or a prevalence for said disease.

The invention furthermore relates to a diagnostic composition comprising the polynucleotide, the gene, the vector, the polypeptide or the antibody of the invention.

In addition, the invention relates to a pharmaceutical composition comprising the polynucleotide, the gene, the vector, the polypeptide or the antibody of the invention. These pharmaceutical compositions comprising, e.g., the antibody may conveniently be administered by any of the routes conventionally used for drug administration, for instance, orally, topically, parenterally or by inhalation. Acceptable salts comprise acetate, methylester, HCl, sulfate, chloride and the like. The compounds may be administered in conventional dosage forms prepared by combining the drugs with standard pharmaceutical carriers according to conventional procedures. These procedures may involve mixing, granulating and compressing or dissolving the ingredients as appropriate to the desired preparation. It will be appreciated that the form and character of the pharmaceutically acceptable character or diluent is dictated by the amount of active ingredient with which it is to be combined, the route of

administration and other well-known variables. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. The pharmaceutical carrier employed may be, for example, either a solid or liquid. Exemplary of solid carriers are lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and the like. Exemplary of liquid carriers are phosphate buffered saline solution, syrup, oil such as peanut oil and olive oil, water, emulsions, various types of wetting agents, sterile solutions and the like. Similarly, the carrier or diluent may include time delay material well known to the art, such as glyceryl mono-stearate or glyceryl distearate alone or with a wax.

The dosage regimen will be determined by the attending physician and other clinical factors; preferably in accordance with any one of the above described methods. As is well known in the medical arts, dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. Progress can be monitored by periodic assessment.

Furthermore, the use of pharmaceutical compositions which comprise antisense-oligonucleotides which specifically hybridize to RNA encoding mutated versions of the polynucleotide or gene according to the invention or which comprise antibodies specifically recognizing a mutated polypeptide of the invention but not or not substantially the functional wild-type form is conceivable in cases in which the concentration of the mutated form in the cells should be reduced.

In another embodiment the present invention relates to the use of the polynucleotide, the gene, the vector, the polypeptide, the polynucleotides having at a position corresponding to position 270 (exon 3) and 206 (exon 8) (GenBank accession No: AF136833.1 and AF136842.1, respectively) of the CYP2C8 gene (GenBank accession No: GI: 13787189) an A instead of a G in position 270 and a G instead of an A in position 206, or the antibody of the invention for the preparation of a diagnostic composition for diagnosing a disease.

Thanks to the present invention the particular drug selection, dosage regimen and corresponding patients to be treated can be determined in accordance with the present invention. The dosing recommendations will be indicated in product labeling

by allowing the prescriber to anticipate dose adjustments depending on the considered patient group, with information that avoids prescribing the wrong drug to the wrong patients or at the wrong dose.

In a further embodiment the present invention relates to the use of the polynucleotide, the gene, the vector, the polypeptide, the polynucleotides having at a position corresponding to position 117 in exon 5 (GenBank accession No: AF136837.1) of the CYP2C8 gene (GenBank accession No: GI: 13787189) a T instead of an A, or the antibody of the invention for the preparation of a pharmaceutical composition for treating a disease.

In a more preferred embodiment of the use of the present invention said disease is an incompatibility or disease related to arachidonic acid metabolism, cancer or cardiovascular disease.

Finally, the present invention relates to a diagnostic kit for detection of a single nucleotide polymorphism comprising the polynucleotide, the gene, the vector, the polypeptide, the antibody, the host cell, the transgenic non-human animal or the solid support of the invention.

The kit of the invention may contain further ingredients such as selection markers and components for selective media suitable for the generation of transgenic cells and animals. The kit of the invention can be used for carrying out a method of the invention and could be, inter alia, employed in a variety of applications, e.g., in the diagnostic field or as research tool. The parts of the kit of the invention can be packaged individually in vials or other appropriate means depending on the respective ingredient or in combination in suitable containers or multicontainer units. Manufacture of the kit follows preferably standard procedures which are known to the person skilled in the art. The kit may be used for methods for detecting expression of a mutant form of the polypeptides, genes or polynucleotides in accordance with any one of the above-described methods of the invention, employing, for example, immunoassay techniques such as radio-immunoassay or enzyme-immunoassay or preferably nucleic acid hybridization and/or amplification techniques such as those

described herein before and in the Examples as well as pharmacokinetic studies when using non-human transgenic animals of the invention.

The figures illustrate the invention:

Figure 1

Correlation of the SNP C104G (Exon 5, I264M) with reduced protein levels of CYP2C8. Expression levels of 14 individuals were determined by Western Blot analysis and LC-MS using verapamil as specific substrate. The boxplots show the distribution of samples according to the genotype at amino acid position 264. The genotype-phenotype correlation is significant ($p = 0.037$, $N=14$).

Figure 2

Correlation of the SNP -370 relative to the start codon ATG with increased expression levels as detected by western blotting, using the drug taxol (paclitaxel) as specific substrate. As shown in the boxplots the genotype-phenotype correlation is significant between homozygous wild type and heterozygous mutant samples ($p=0.044$, $N=20$). (One homozygous sample was analyzed and yielded expression levels > 400).

Figure 3

Correlation of the SNP at position -370 in the untranslated region based on deviates from mean values of CYP2C8 expression levels from two independent sample collectives ($N = 62$). The phenotype of the patients was determined by LC/MS and Western Blot. The phenotype-genotype correlation is significant as shown in the boxplots ($p = 0.017$ for homozygous vs wildtype and $p = 0.071$ for heterozygous vs wildtype).

Figure 4

Correlation of the allele including the linked SNPs G-1207A, delAT-640/41 (both in the promoter), G270A (exon 3), and A206G (exon 8) with a poor metabolizer phenotype (PM). However, no homozygous individuals for this allele could yet be phenotyped. The data shown in the box plot do only show a trend ($p = 0,071$; $N = 18$) but no significance.

Figure 5

Transfection of LS174T cells with either a CYP2C8 wild type promoter construct or two SNP(s) containing promoter fragments (G-1207A and -640 to 641delAT or T-370G). Using the eukaryotic pGL3 expression vector mean values of six independent transfection assays were analysed following normalisation of 2C8 wild type activity to 100%.

Figure 6

Computer modeled CYP2C8 enzyme structure of the protein variants Thr 159 Pro (frameshift), Glu 274 Stop and Gly 365 Ser. Dark grey represents the unchanged structure of the variant protein, the light grey prepresents the missing amino acids of the CYP2C8 variant structure. The circle indicates the active site of the enzyme with the altered amino acid Gly 365 Ser in dark.

Figure 7

Reference or wild type GenBank sequences for the polynucleotides, polypeptide and mRNA according to the present invention

The invention will now be described by reference to the following biological Examples which are merely illustrative and are not constructed as a limitation of the scope of the present invention.

Examples

Example 1: Isolation of genomic DNA from human blood, generation and purification of CYP2C8 gene fragments

Genomic DNA was obtained by standard ion exchange chromatography techniques (Qiagen kits for isolation of genomic DNA from blood). Blood from all the individuals tested (volunteers from Parexel, Berlin and the Institute for Clinical Pharmacology, Stuttgart) was obtained under consideration of all legal, medical and bureaucratic requirements. Further samples from other ethnic populations (e.g. Caucasian, Japanese, African-American) were purchased from commercial sources.

By using polymerase chain reaction (PCR) with specific oligonucleotide primers, two for each fragment, defined DNA-fragments containing specific parts of the human CYP2C8 gene were obtained. These specific oligonucleotide primers were designed to bind to sequences upstream and downstream of the various exons as well as in the 5 prime region of the CYP2C8 gene. The resulting DNA fragments did not contain codogenic parts alone but also sequences covering the intronic parts located at the exon-intron boundaries. These sites are known to be important for correct processing and subsequent expression of the protein encoding mRNA, a process called „splicing“. Commercially synthesized oligonucleotide primer pairs that were purified by affinity chromatography were optimized for each of the 9 exon and 4 promoter fragments of the human CYP2C8 gene. The sequences for each primer are listed in table 1.

Polymerase chain reactions were performed under conditions that were optimized for each of the nine fragments and a promoter region covering about 2kb upstream of the initiation codon for mRNA translation. PCRs were carried out for all exons in a volume of 50 µl. 10 – 50 ng of template DNA were added to standard PCR-buffer

containing 1,5 mM MgCl₂, 200 µM dNTPs and 1 U Taq-polymerase (all from Qiagen, Hilden) as well as 10 - 40 pMol of primers (MWG Biotech, Munich). All PCR reactions were performed on identical conditions at a Perkin Elmer thermocycler (Modell 9700) with an initial denaturation step of 94°C for 2 min, followed by 34 cycles for PCR-fragment generation with 45 s denaturation at 94°C, 45 s of annealing at 62°C and 1 min at 72°C for elongation. The exact location of the primers and size of the desired fragments are also listed in table 1.

The defined DNA fragments containing specific parts of the CYP2C8 gene, exon as well as some intron sequences at the intron-exon boundaries were processed to remove nonincorporated nucleotides and buffer components that might otherwise interfere with the subsequent determination of the individual CYP2C8 genotype by direct cycle sequencing. For this purification, standard ion exchange chromatography techniques were used (Qiagen kits for PCR-fragment purification). For all fragments sufficient yields of purified fragments, suitable for direct DNA sequencing analysis were obtained. Aliquots of purified fragments were subjected to direct sequence analysis of the CYP2C8 gene in an ABI 3700 capillary sequencer.

Example 2: Identification of different CYP2C8 alleles by sequence determination in various individuals

For sequence analysis of relevant regions of the human CYP2C8 gene from many different individuals, PCR amplifications of the relevant regions of the gene were carried out (primers see table 1) following purification of the PCR products and sequencing with established methods (ABI dye terminator cycle sequencing). Since the individual genetic makeup is represented by two copies of any gene (diploidy), great care has to be taken in the evaluation of the sequences not to unambiguously identify homozygous, but also heterozygous sequence variation. Therefore, in cases where no clear discrimination could be detected, forward and reverse sequencing was performed. Moreover, for the discovery of complete and defined alleles, e.g. in linkage equilibrium, it is necessary to cover all exons as well as the promoter region to provide a comprehensive basis for the phenotype prediction of individual SNPs.

For the evaluation of CYP2C8 variations in the human population, sequence analyses of the relevant regions, including a 2kb promoter fragment and all exons of the gene were carried out from genomic DNA of each 48 Caucasian, Japanese and African-American individuals. The sequences were subjected to a computer analysis programme (PhredphrapTM, Perkin Elmer) and inspected manually for the occurrence of DNA sequences deviating from recently published CYP2C8 sequences that were considered to represent the „wild type“ sequences in this work.

Because population genetics enables a calculation of the expected frequency of homozygous vs. heterozygous alleles of a defined gene (Hardy Weinberg formula: $2p^2 + 2pq + 2q^2 = 1$), it was possible to confirm predicted distributions of homozygous vs. heterozygous alleles and deviations from the experimental findings. This serves as experimental control that a detected sequence variation indeed represents a novel allele.

Several new CYP2C8 sequence variations were discovered and experimentally confirmed using this approach which are shown in table 2. 18 polymorphisms are located in the 5' untranslated region/promoter of the gene (GenBank accession No: AF136830.1). 22 new polymorphisms could be found in sequences of introns 1, 2, 3, 4 and 8 (GenBank accession Nos: AF136832.1, AF136833.1, AF136835.1, AF136843.1 and AF136844.1) and one in the 3' untranslated region of the gene (GenBank accession No: AF136845.1).

Furthermore, three particular nucleotide changes in exon 3 (position 334 at exon /intron boundary, GenBank accession No: AF136833.1) and exon 8 (position 30 and 87, GenBank accession No: AF136843.1) were detected that do not change the amino acid sequence. In exon 3 (position 329, GenBank accession No: AF136833.1), exon 4 (position 243, GenBank accession No: AF136842.2 and position 13 of GenBank accession No: AF136835.1), exon 5 (position 42, 101 and 104, GenBank accession No: AF136837.1), exon 6 (position 309, GenBank accession No: AF136838.1) and 7 (position 1135, GenBank accession No: NM_000770.1 and position 232, GenBank accession No: AF136840.1) nine SNPs could be identified that change the protein sequence as shown in table 4, where the deviative amino acid is typed in a bold style. These novel, and already published CYP2C8 SNPs serve as markers for the characterization of the CYP2C8 status in patients.

The positions of the novel CYP2C8 SNPs, including the exact novel sequence context are listed in table 2. The deviative base in the sequence is typed underlined and in a bold style.

Example 3: Determination of the CYP2C8 promoter allele containing G-1207A, delAT -640 to -641 as a pharmacogenetic factor influencing drug levels

The anticancer drug taxol (paclitaxel) can be considered to be the prototype for CYP2C8 and it's isoform 6-hydroxypaclitaxel as diagnostic substrate. Furthermore, verapamil represents another specific substrate for CYP2C8 that is methylated to different metabolites, e.g D-702 and D-703 or desalkylated to D-617 (i.e. a substrate of CYP3A4). The generation of a specific metabolite used to selectively proof the functional acitvity of CYP2C8 (LC-MS) is shown in figure 4. In parallel, the amount of enzyme is determined by western blotting (see example 9).

To validate a possible correlation of the single nucleotide polymorphisms G-1207A, delAT -640 to -641, and/or T-370G (all 5'UTR/Promoter) and/or C104G (exon 5) with a certain phenotype, biopsies from patients of two different studies were analyzed. The functional characterization of the CYP2C8 gene had been determined from enterocyte preparations of the duodenum and the liver.

Using these analytical tools, phenotypically characterized samples that have been treated with taxol (N = 22) or verapamil (N = 15, 44) were subjected to genotyping. Either of the collectives showed that two SNPs G-1207A, delAT-640 to -641 of the 5' untranslated region are in linkage disequilibrium and, in combination with SNPs G270A (exon 3) and A206G (exon 8), represent a new allele (haplotype) that is mainly defined by the presence of the two novel promoter SNPs G-1207A and delAT-640 to -641. Figure 4 shows that this allele is responsible for the reduced level of metabolized substrate.

Example 4: Functional consequences of the identification of CYP2C8 promoter polymorphisms

The eukaryotic promoter region of a gene is composed of several regulatory elements, e.g. enhancer, silencer and other responsive elements. Here, single nucleotide polymorphisms exhibit significant influence. Regarding cytochrome P450-enzymes induction mechanisms, e.g. transcription factors like C/EBP, HPFs or barbiebox-sites identified in CYP2C9 (Klose, J Biochem Mol Toxicol 13 (1999), 289-95) are important since temporary expression is required. SNPs change or interfere with such elements and can alter promoter action and/or transcription activity. The novel SNP identified at position -1207 most probably abolishes transcription factor binding that has selectively been shown for the binding of tissue specific sterol regulatory element binding protein 1 (SREBP-1, Vallett, J Biol Chem. 271 (1996), 12247-53). Reduced expression has been shown if position -1207 does not correspond to the wild type. The correlation of this new allele with CYP2C8 enzyme activity is displayed in figure 4. The linkage with another unidentified SNPs further downstream in the promoter region of CYP2C8 (delAT -640 to -641) confers new value to the allele in respect to diagnostic applications.

Example 5: Determination of the CYP2C8 promoter polymorphism T-370G as a pharmacogenetic factor influencing drug levels

Another polymorphism located further downstream at position 1668 in the 5-prime untranslated region (position -370 relative to the ATG-start codon) could be identified to significantly increase the expression level as shown in figure 3. This allele refers to as an extensive metabolizing phenotype (EM) that was confirmed by investigation of phenotypically characterized samples. In several cases, single nucleotide polymorphisms G-1207A, delAT -640 to -641 occur in combination with T-370G. LC-MS results from these samples show that individuals carrying the T-370G alone have an increased CYP2C8-activity as compared to those heterozygous for the polymorphisms G-1207A, delAT -640 to -641 allele. Concerning the application in a diagnostic assay these data clearly show the influence of position T-370G on the expression levels of CYP2C8, i.e. the latter SNP is responsible for a change from poor (PM) to intermediate/extensive metabolism (IM/EM) as demonstrated in example 9/table 6.

Independently, a further validation was carried out by genotyping 22 samples corresponding to individual liver extracts, in which the metabolism of taxol was assessed. The data confirm results as presented in examples 8 and 9. Figure 2 shows the significant correlation between a heterogeneous polymorphism at position -370G and the increase of CYP2C8 protein levels. The liver extract from a sample homogeneous for position -370 displayed the highest CYP2C8-protein level (>400pmol/mg protein). This is an additional independent result that supports the significant correlation in figure 2.

Example 6: Determination of the CYP2C8 polymorphism at amino acid position 264 (exon 5) as a pharmacogenetic factor influencing drug levels

In another embodiment the present invention relates to a polymorphism in exon 5 at position C104G (GenBank accession No: AF136837.1). This change correlates with a reduced protein concentration analyzed from genotyped samples (figure 1) that could be due to less stable mRNA or protein. The polypeptides encoded by the polynucleotides of the invention may have altered biological or immunological properties due to the polymorphisms referred to in accordance with the present invention. Examples for said altered properties are stability of the polypeptides which may be effected or the incapability to effectively metabolize certain drugs.

Example 7: Using Restriction Fragment Length Polymorphism (RFLP) analysis to detect SNPs relevant in phenotypic prediction

CYP2C8-polymorphisms can be detected not only by sequencing but also by various other means. As one alternative to the sequencing methodology, genotyping can be performed with PCR fragments to be processed by one restriction endonuclease specifically cutting at a region, composed of a unique sequence of 4-6 nucleotides. Due to the limited length of a PCR-fragment sometimes advantage can be taken of this specificity if it discriminates mutant and wild type, i.e. resulting in digested or undigested fragments (Table 5). Regarding the present invention this was the case for

fragments of the promoter region 4 (position -370, double cutter AcsI for wild type and single cutter in the mutant), that indicates the respective allele, exon 3 (position 275, single cutter SapI for CYP2C8-mutant), and exon 5 (position 104, single cutter ClaI for wild type). Depending of the fragments' specific SNP-region the restriction pattern unambiguously reflects either the wild type, the heterozygous or homozygous mutant. As defined by the primers listed in table 1, the exons screened for result in the following RFLP-fragments:

Table 5:

SNP position (PCR-fragment)	Enzyme	Length (bp)	Genotype		
			wt/wt (bp)	wt/mut (bp)	mut/mut (bp)
# -370, (5'UTR)	AcsI	483	33/150/ 300	33/150/183/300	183/300
# 270, (exon 3)	SapI	328	328	149/179/328	149/179
# 104, (exon 5)	ClaI	584	192/392	192/392/584	584

Example 7: Identification of new CYP2C8 polymorphisms by sequence analysis of a collection of various individuals from different ethnic groups

The screen for SNPs in the CYP2C8 gene in the genomes of different ethnic groups yielded a number of polymorphisms listed in table 2. 48 samples were analyzed from each of the ethnic populations Caucasian, Japanese and African-American, respectively. Within this collection, the large number of SNPs in the untranslated region could be considered to potentially influence the protein level.

Furthermore, several polymorphisms show extensive inter ethnic discrepancies (Table 3) between various ethnical groups. The 57 new polymorphisms identified in the CYP2C8 gene will complement the existing knowledge and contribute to a more comprehensive understanding of the gene, avoiding problems in drug response and, concerning other ethnical groups, thus facilitating "bridging studies" that could be of interest when projecting data from these embodiment.

Example 8: Characterization of promoter SNPs by transfection of promoter- reporter plasmids into human cells

Three promoter SNPs were tested for their contribution to different expression levels by transfection assays using the LS174T cell line. Promoter fragments containing the wild type or the SNPs at positions corresponding to positions G-1207A and delAT - 640 to - 641 (GenBank accession No: AF136830.1) or at position corresponding to position T-370G (GenBank accession No: AF136830.1) were introduced into a commercial mammalian expression vector. The plasmid harbours standard sequences for the propagation in eukaryotic cells including the reporter gene *luciferase* that is controlled by the integrated promoter sequence. Following sequence verification of each DNA-insert, cells were cotransfected with β -galactosidase, harvested after 48h and analysed for luciferase activity. Promoter activities (%) are shown in figure 5 following normalization to the transfection efficacy as determined by β -galactosidase detection. The data are in full agreement with the observations from phenotypically characterized samples. A DNA-construct that contains the 1207G>A and -640 to -641 delAT polymorphisms (figure 5) showed decreased transcription for the CYP2C8- promoter levels compared to the wild type. In contrast, the reporter plasmid revealed increasing levels of luciferase protein under control of a CYP2C8-promoter containing a polymorphism corresponding to position - 370.

Example 9: Protein quantification of samples containing SNPs at promoter position G-1207A, delAT - 640 to - 641, T-370G and C104G (exon 5, amino acid position 264) of CYP2C8

Protein extracts have been prepared from human liver samples. The protein levels of CYP2C8 were analysed by western blot using samples genotyped for SNPs G-1207A, delAT -640 to -641, T-370G (all GenBank accession No: AF136830.1) and C104G (GenBank accession No: AF136837.1). Table 5 shows the effects of different genotypes on the expression levels of CYP2C8 (pmol/mg) normalized for the wild type (=100%). Results are in total agreement with the functional data described by

the promoter-reporter assays in example 8. The promoter SNP in position T-370G confers to increased levels of the CYP2C8 protein (150%), whereas polymorphisms G-1207A and delAT -640 to -641 in contrast show a reduced protein expression (72%). In combination with SNPs G-1207A and delAT -640 to -641, or C104G alone the polymorphism T-370G differentially influences protein levels as indicated by arrow. Here, the presence of two SNPs with significant frequency leads to combined effects. Therefore, considerations for reliable phenotype prediction as a result from genotyping must depend on multiple SNP-analyses. The data indicate that the SNP in position -370, which by itself is responsible for up regulation of the CYP2C8-protein level, shows no significant CYP2C8 increase if it is combined with SNPs G-1207A and delAT -640 to -641 (5'UTR), which by themselves reduce the expression. The combined expression level of 119% is barely higher than in the homozygous wildtype situation. Vice versa, in combination with the C104G allele (exon 5) the increased expression due to the SNP at position -370 is compensated by the SNP C104G to normal expression levels (97%) compared to the wild type. This reflects the strong impact of SNP C104G alone on the protein level (see figure 1). The SNP C104G therefore represents an allele for down regulation.

Table 6:

Genotype (by detection of listed SNPs)	CYP 2C8-levels (%)	Effects
Wild type	100	No
G-1207A, delAT -640 to -641 (5'UTR)	72	↓
T-370G (5'UTR)	150	↑
G-1207A, delAT -640 to -641 and T-370G (5'UTR)	119	↑↓
T-370G (5'UTR) and aa I264M (exon 5)	97	↑↓

No sample with aa change I264M or G-1207A, delAT -640 to -641 (5'UTR) and I264M was detected.

Example 10: Pharmacogenetic relevance of the CYP2C8 polymorphisms at position 329 (exon 3, Thr 159 Pro), 309 (exon 6, Glu 274 stop) and 1135 (exon 7, Gly 365 Ser)

In another embodiment the present invention relates to a polymorphisms in exon 3, exon 6 and 7 at positions 329 (GenBank accession No: AF136833.1), 309 (GenBank accession No: AF136838.1) and 1135 (GenBank accession No: NM_000770.1) respectively. The delA change in position 329 causes a frameshift abolishing the C-terminal part of the protein. The G309T change in position 309 results in a premature termination at amino acid position 274 of the protein. Both variant transcripts encode for polypeptides that will lose their function and are therefore most likely poor metabolizer alleles (Fig. 6). In another embodiment the present invention relates to a polymorphism at position G1135A in exon 7 (GenBank accession No: NM_000770.1). This substitution results in a change from Glycine to Serine at position 365 within the active site of the CYP2C8 enzyme (Fig. 6). Because the active site of wildtype CYP2C8 contains hydrophobic amino acids to enable the hydrophobic substrate to efficiently enter the substrate pocket, this amino acid exchange to a hydrophilic residue will severely interfere with substrate binding and subsequent metabolism.

Table 1 Primer sequences for the generation of CYP2C8 PCR-fragments

All primer locations refer to different contigs of HTGS-Database, GenBank Acc. No. AL359672.10
 PCR-fragments at the 5'UTR are overlapping.

PCR-fragment name	PCR- fragm. size (bp)	Contig spec. exon location	Primer position	Primer Sequence (5' - 3')
Contig		115244 - 120972 (5629bp)		
5'UTR Fragm. 1	537	534	120439 - 120416 forward: 119902 - 119926 reverse:	ATT TTA GTC AAT CTT GGT GGC CCG TTC AAC AGA AGA TGG AAC ACA GGG A
5'UTR Fragm. 2	545		120004 - 119980 forward: 119460 - 119483 reverse:	TCA TGA CCA TTG ACT ATC AGT TCC C TGA TAC CCA TTG GGG TTC ATT ACC
5'UTR Fragm. 3	751		119576 - 119552 forward: 118826 - 118854 reverse:	AAC AGA GTC AAG GTG GCG TAT CTT C CAA TAT TCT CAG ATT AAT GAC CAG TTG GG
Sequencing primer			118938 - 118963	AGA CTT AGC CCT TGA TAA CAA AAG CC
5'UTR Fragm. 4	483	-2486	119008 - 118982 forward: 118526 - 118550 reverse:	GTT TAG GCA GCT GTA TTT TAA GTG AAC ACT CCA AAG TTT TTA TAA CAC TCC C
Exon 1	472	2487 - 2654	118687 - 118664 forward: 118216 - 118242 reverse:	GGC ACT GGA AAG AAG GAG TAG GAC GAT CTA TTA TAA TAG TGT GCT TCC AGG
Exon 2	457	4198 - 4360	116999 - 116978 forward:	TTG TGT ACC AAT TGC CTG GGT C

Exon 3	328	4532 - 4681	116543 - 116566	reverse:	TTT TTA GGG CTC TGT TTT CCA TCC
Contig		78619 - 85206 (6588 bp)			
Exon 4 Alternative	541	1378 - 1538 AF136834.2	83947 - 83970 44-67	forward:	TCC ATG CTG ATT TTT TTT GGA CAC
Contig		138518 - 143654 (5137bp)	83429 - 83450	reverse:	CTG ACC CCT TGC ACT TCT GAT G
Exon 5	583	1319 - 1495	139593 - 139617 140176 - 140154	forward: reverse:	TGA CGA GTT ATT GGG TGC AGT ACA C TTC CAT GAT GTT TAG TGC TGC AGG CC
Contig		85307 - 115243 (29937bp)			
Exon 6	519	8875 - 9016	93884 - 93903 94402 - 93379	forward: reverse:	TTG AAG TAA GAC AGG GCA TCG G AGA AAC AAG GTG GAG GAT ACT GGC
Exon 7	328	11749 - 11936	96986 - 97009 97313 - 97290	forward: reverse:	GGC CAT GAA TTG CTA TGA CAA ATG GGT TGG AAC CAA ACC AGC ACT ATG
Exon 8	462	15788 - 15929	100976 - 100997 101437 - 101418	forward: reverse:	CTG GCT GGA CCT GAG TTT CCT C TTA ACT CCT GCA AGC CCC GC
Exon 9/3'UTR	543	17526 - 17707	102630 - 102652 103172 - 103149	forward: reverse:	GTA CAT TTG TTT GTC CCA CCA TCC TGC AGT GAC CTG AAC AAC TCT CCT

Table 2 SNPs identified in the CYP2C8 gene

PCR- fragm Location	GenBank Acc. No	variant position (relative to ATG)	wild type (f) and (r)	wild type/mutant (f) and (r)	mutant/mutant (f) and (r)
5'UTR- fragm 1	AF136830.1	#306 to307 (= -1731 to -1732)	f : GATGTGATG <u>A</u> GTGTGAAAAAT r : ATTTTCACA <u>C</u> TCAATCACATC	f : GATGTGATG (<u>AG</u>)TGTGAAAAAT f : GATGTGATGTGTGAAAAAT r : ATTTTCACA (<u>CT</u>)CATCACATC r : ATTTTCACACATCACATC	
5'UTR- fragm 1	AF136830.1	#411 (= -1627)	f : GGAAATAA <u>A</u> CT <u>T</u> AGTACTGGTC r : GACCAGTAC <u>A</u> <u>T</u> GTATTATTCC	f : GGAAATAA <u>A</u> CT/ <u>AG</u> TACTGGTC f : GGAAATAA <u>A</u> CTACTGGT(r : GACCAGTAC <u>A</u> / <u>T</u> GTATTATTCC r : GACCAGTAC <u>T</u> GTATTATTCC	⁴⁸
5'UTR- fragm 1	AF136830.1	#560 (= -1478)	f : GGTCTGCAC <u>A</u> TTTGCAGTGG r : CCACTGCAA <u>T</u> GTGCAGACC	f : GGTCTGCAC <u>A</u> / <u>G</u> TTGCAGTGG f : GGTCTGCAC <u>G</u> TTTGCAGTGG r : CCACTGCAA <u>C</u> / <u>T</u> GTGCAGACC r : CCACTGCAA <u>C</u> GTGCAGACC	
5'UTR- fragm 2	AF136830.1	#713 (= -1325)	f : AAAACAATA <u>G</u> AAGCAGCCA r : TGGCTGCTT <u>C</u> TATTGTTTT	f : AAAACAATA <u>G</u> / <u>T</u> AAGCAGCCA f : AAAACAATA <u>T</u> AAGCAGCCA r : TGGCTGCTT <u>A</u> / <u>C</u> TATTGTTTT r : TGGCTGCTT <u>A</u> TATTGTTTT	

5'UTR- fragm 2	AF136830.1	#817 (= - 1221)	f : AGTGTCTGAAC <u>C</u> AACTTTTCAC r : GTGAAAGTT <u>G</u> TTTCAGCACT	f : AGTGTCTGAAC <u>A</u> AACTTTTCAC r : GTGAAAGTT <u>T</u> TTTCAGCACT
5'UTR- fragm 2	AF136830.1	#824 (= - 1214)	f : AACAACTTT <u>C</u> ACTTGTGAG r : CTCACAAAGT <u>G</u> AAAGTTGTT	f : AACAACTTT <u>A</u> ACTTGTGAG r : CTCACAAAGT <u>T</u> AAAGTTGTT
5'UTR- fragm 2	AF136830.1	#831 (= - 1207)	f : TTCACCTTGT <u>G</u> AGGTGATGC r : GCATCACCT <u>T</u> CACAAAGTGAA	f : TTCACCTTGT <u>A</u> AGGTGATGC r : GCATCACCT <u>T</u> ACAAAGTGAA
5'UTR- fragm 2	AF136830.1	#879 (= - 1159)	f : CTTTTTGAGC <u>G</u> TCTCCGGTC r : GACCCGGAGAG <u>C</u> GCTCAAAAG	f : CTTTTTGAGC <u>A</u> TCTCCGGTC r : GACCCGGAGAG <u>T</u> GCTCAAAAG
5'UTR- fragm 2	AF136830.1	886 (= - 1152)	f : GCGTCTCCG <u>G</u> TCTCTTAT r : ATAAGAGGA <u>C</u> CGGAGACGC	f : GCGTCTCCG <u>T</u> TCTCTTAT r : ATAAGAGGA <u>A</u> CGGAGACGC
5'UTR- fragm 3	AF136830.1	#1058 (= - 980)	f : ACCCCAATG <u>G</u> AGTATCAGAA r : TTCTGATAC <u>C</u> CAATTGGGGT	f : ACCCCAATG <u>A</u> AGTATCAGAA r : TTCTGATAC <u>T</u> CAATTGGGGT
5'UTR- fragm 3	AF136830.1	#1271 to 1273	f : GTATTTATG <u>T</u> TAATTATGT r : GTATTTATG(<u>TTA</u>)TTAATTATGT	f : GTATTTATG(<u>TTA</u>)TTAATTATGT r : GTATTTATGTTAATTATGT

	(= - 765) to (- 767)	r: ACATAATAA <u>TAA</u> CATAAATAC	r: ACATAATAA (<u>TAA</u>) CATAAATA r: ACATAATAACATAAAATAC C
5'UTR- fragm 3	AF136830.1 #1397 to 1398 (= - 640) to (- 641)	f: TGTAAATAAC <u>AT</u> ATATATTTA r: TAAATATATAT <u>AT</u> GTTATTACA	f: TGTAAATAAC (<u>AT</u>) ATATATTTA f: TGTAAATAACATATATTTA r: TAAATATATAT (<u>AT</u>) GTTATTACA r: TAAATATATGTTATTACA
5'UTR- fragm 4	AF136830.1 #1627 (= - 411)	f: TTTTTTTATAT <u>T</u> ACAAAAATAT r: ATATTTTGT <u>A</u> TATATAAAAA	f: TTTTTTTATAT <u>T</u> /CACAAAAATAT f: TTTTTTTATAT <u>C</u> ACAAAAATAT r: ATATTTTGT <u>G</u> / <u>A</u> TATATAAAAA r: ATATTTTGT <u>G</u> TATATAAAAA
5'UTR- fragm 4	AF136830.1 #1668 (= - 370)	f: GGTCATAAA <u>T</u> TCCCAACTG r: CAGTTGGGA <u>A</u> TTTATGACC	f: GGTCATAAA <u>T</u> /GTCCTCAACTG f: GGTCATAAA <u>G</u> TCCCAACTG r: CAGTTGGGA <u>C</u> / <u>A</u> TTTATGACC r: CAGTTGGGA <u>C</u> TTTATGACC <u>S</u>
5'UTR- fragm 4	AF136830.1 #1767 (= - 271)	f: ACATTTGGAA <u>C</u> AAACCAGGGA r: TCCCTGGTT <u>G</u> TTCCCAATGT	f: ACATTTGGAA <u>C</u> / <u>A</u> AAACCAGGGA f: ACATTTGGAA <u>A</u> AAACCAGGGA r: TCCCTGGTT <u>T</u> / <u>G</u> TTCCCAATGT r: TCCCTGGTT <u>T</u> TTCCCAATGT
5'UTR- fragm 4	AF136830.1 #1785/ 1786 (= - 252)	f: AATTAATAAAATACCTGGGC r: GCCCAGGTATTTTAAATT	f: AATTAATAAAA (<u>A</u>) TACCTGGGC f: AATTAATAAAA <u>A</u> TACCTGGGC r: GCCCAGGTA (<u>T</u>) TTTTAAATT r: GCCCAGGTA <u>T</u> TTTTTAAATT
5'UTR- fragm 4	AF136830.1 #1887 (= - 151)	f: CTATCCATG <u>G</u> GGCCAAAAGTC r: GACTTTGGC <u>C</u> CCATGGATAG	f: CTATCCATG <u>G</u> / <u>A</u> GGCCAAAAGTC f: CTATCCATG <u>A</u> GGCCAAAAGTC r: GACTTTGGC <u>T</u> / <u>C</u> CATGGATAG r: GACTTTGGC <u>T</u> CATGGATAG
5'UTR- fragm 4	AF136830.1 #1905 (= - 133)	f: CCACTCAGAA <u>A</u> AAAAAGTAT	f: CCACTCAGAA/ <u>C</u> AAAAAGTAT f: CCACTCAGAA <u>C</u> AAAAAGTAT

5'UTR- fragm 4	AF136830.1	#1952 (= - 86)	r: ATACTTTT <u>T</u> TCTGAGTGG	r: ATACTTTT <u>T</u> TCTGAGTGG	r: ATACTTTT <u>T</u> TCTGAGTGG
			f: ACATGTCAA <u>A</u> AGAGACACAC	f: ACATGTCAA <u>A</u> CGAGACACAC	f: ACATGTCAA <u>C</u> CGAGACACAC
			r: GTGTGTCTC <u>T</u> TTGACATGT	r: GTGTGTCTC <u>G</u> /T <u>T</u> TGACATGT	r: GTGTGTCTC <u>G</u> T <u>T</u> TGACATGT
Intron 1	AF136832.1	#171	f: ATTCAGAAA <u>T</u> ATCGAATCT	f: ATTCAGAAA <u>T</u> CATCGAATCT	f: ATTCAGAAA <u>C</u> ATCGAATCT
			r: AGATTGAT <u>A</u> TTTCTGAAT	r: AGATTGAT <u>G</u> /A <u>T</u> TTCTGAAT	r: AGATTGAT <u>G</u> T <u>T</u> TTCTGAAT
Intron 1	AF136832.1	#258	f: AGCAAAATAG <u>C</u> GGACTTATTT	f: AGCAAAATAG <u>C</u> /TGACTTATTT	f: AGCAAAATAG <u>T</u> GGACTTATTT
			r: AAATAAGTC <u>G</u> CTATTTGCT	r: AAATAAGTC <u>A</u> /GCTATTTGCT	r: AAATAAGTC <u>A</u> CTATTTGCT
Intron 2	AF136833.1	#122	f: ATGGCTGCC <u>G</u> AGTGTGCA	f: ATGGCTGCC <u>G</u> /AAGTGTGCA	f: ATGGCTGCC <u>A</u> AGTGTGCA
			r: TGCAACACT <u>C</u> GGCAGCCAT	r: TGCAACACT <u>T</u> /CGGCAGCCAT	r: TGCAACACT <u>T</u> TGGCAGCCAT ²
Intron 2	AF136833.1	#150	f: TCCTTGGCT <u>G</u> TGAATTCTC	f: TCCTTGGCT <u>G</u> /ATGAATTCTC	f: TCCTTGGCT <u>A</u> TGAATTCTC
			r: GAGAAATTC <u>A</u> CAGCCAAAGGA	r: GAGAAATTC <u>A</u> T/CAGCCAAAGGA	r: GAGAAATTC <u>A</u> TAGCCAAAGGA
Intron 2	AF136833.1	#180/181	f: CCTTTTTTTTATTAGGAAT	f: CCTTTTTTTT(<u>T</u>)ATTAGGAAT	f: CCTTTTTTTT <u>T</u> ATTAGGAAT
			r: ATTCCCTAATAAAAAAAG	r: ATTCCCTAAT(<u>A</u>)AAAAAAGG	r: ATTCCCTAAT <u>A</u> AAAAAAGG
Intron 2	AF136833.1	#182	f: CTTTTTTTATATAGGAATCA	f: CTTTTTTTAT/CTAGGAATCA	f: CTTTTTTTAT <u>C</u> TAGGAATCA
			r: TGATTCCCTA <u>A</u> TAAAAAAG	r: TGATTCCCTAG/A <u>T</u> AAAAAAG	r: TGATTCCCTAG <u>T</u> AAAAAAG
Exon 3	AF136833.1	#270	f: TGGGGAAGAGGAGCATTGA	f: TGGGGAAGAG/ <u>A</u> GAGCATTGA	f: TGGGGAAGAA <u>A</u> GAGCATTGA
			r: TCAATGCTC <u>C</u> TCTTCCCCA	r: TCAATGCTC <u>T</u> /CTCTTCCCCA	r: TCAATGCTC <u>T</u> TCTTCCCCA
Exon 3	AF136833.1	#334	f: AAAAACCAAGGGTGGGTGA	f: AAAAACCAAG/ <u>A</u> GGTGGGTGA	f: AAAAACCAAA <u>A</u> GGTGGGTGA
			r: TCACCCACC <u>C</u> TTGGTTTTT	r: TCACCCACC <u>T</u> /CTTGGTTTTT	r: TCACCCACC <u>T</u> TTGGTTTTT

Exon 3	AF136833.1	#329	f: TTGAGAAAAA <u>A</u> CCAAAGGGTG r: CACCCCTGGT <u>T</u> TTTTCTCAA	f: TTGAGAAAAA <u>(A)</u> CCAAAGGGTG r: CACCCCTGGT <u>(T)</u> TTTTCTCAA	f: TTGAGAAAAA <u>C</u> CAAGGGTG r: CACCCCTGGT <u>T</u> TTTTCTCAA
Intron 3	AF136833.1	#378	f: CAGTTACCT <u>G</u> TCTTCACTA r: TAGTGAAGAC <u>C</u> AGGTAACTG	f: CAGTTACCT <u>G</u> /C <u>T</u> TCTCACTA r: TAGTGAAGAG/CAGGTAACTG	f: CAGTTACCT <u>C</u> TCTTCACTA r: TAGTGAAGAGAGGTAACTG
Intron 3	AF136834.2	#87	f: TGTAAGATA <u>T</u> GTTTTAAAT r: ATTTTAAAC <u>A</u> TATCTTACA	f: TGTAAGATA <u>(T)</u> GTTTTAAAT r: ATTTTAAAC <u>(A)</u> TATCTTACA	f: TGTAAGATA <u>T</u> GTTTTAAAT r: ATTTTAAAC <u>T</u> ATCTTACA
Intron 3	AF136834.2	#162	f: ATAATTTT <u>T</u> TAAAAATTT r: AAAATTTTAA <u>A</u> AAAAATTAT	f: ATAATTTT <u>T</u> /A <u>T</u> AAAAATTT r: AAAATTTTAT/A <u>A</u> AAAAATTAT	f: ATAATTTT <u>T</u> TAAAAATTT r: AAAATTTTAT <u>A</u> AAAAATTAT
Intron 3	AF136834.2	#163	f: TAAATTTT <u>T</u> TAAAAATTTT r: AAAATTTT <u>A</u> AAAAAATTA	f: TAAATTTT <u>T</u> /A <u>A</u> AAAAATTTT r: AAAATTTT <u>T</u> /A <u>A</u> AAAAAATTA	f: TAAATTTT <u>T</u> TAAAAATTTT r: AAAATTTT <u>T</u> TAAAAAATTT
Exon 4	AF136834.2 NM_000770.1	#243 #583	f: ATCTGCTCC <u>G</u> TGTGTTTCC r: GGAAAAACA <u>C</u> GGAGCAGAT	f: ATCTGCTCC <u>G</u> /A <u>T</u> TGTTTCC r: GGAAAAACA <u>T</u> /CGGAGCAGAT	f: ATCTGCTCC <u>A</u> TTGTTTCC r: GGAAAAACA <u>T</u> GGAGCAGAT
Exon 4	AF136835.1	#13	f: GGATTCTGA <u>A</u> CTCCCCATG r: CATGGGGAG <u>T</u> TCAGAAATCC	f: GGATTCTGA <u>A</u> /GCTCCCCATG r: CATGGGGAGC/ <u>T</u> TCAGAAATCC	f: GGATTCTGA <u>G</u> CTCCCCATG r: CATGGGGAG <u>C</u> TCAGAAATCC
Intron 4	AF136835.1	#180	f: TGATTTCTC <u>T</u> GTTCAAAATT r: AAATTTGAAC <u>C</u> AGGAAATCA	f: TGATTTCTC <u>T</u> G/A <u>T</u> TCAAAATT r: AAATTTGAAT/ <u>C</u> AGGAAATCA	f: TGATTTCTC <u>T</u> AATTCAAAATT r: AAATTTGAAT <u>T</u> AGGAAATCA
Intron 4	AF136836.1	#116	f: ACTTAAAGT <u>A</u> TAAATAAAA r: TTTTATTAT <u>T</u> ACTTTAAGT	f: ACTTAAAGT <u>A</u> /GTAATAAAAA r: TTTTATTAT <u>C</u> /TACTTTAAGT	f: ACTTAAAGT <u>G</u> TAAATAAAAA r: TTTTATTAT <u>A</u> CACCTTTAAGT

Intron 4	AF136836.1	#132	f: AAAATGTATATATATGTATAA r: TTATACATAATATACATTTT	f: AAAATGTATGTATGTATAA r: TTATACATACTATACATTTT
Intron 4	AF136836.1	#172	f: ATGATGTCTTATTCATATT r: AATATGAATAAGACATCAT	f: ATGATGTCTCATTCATATT r: AATATGAATGAGACATCAT
Intron 4	AF136836.1	#189	f: TTTATAGTTATAATTTCAA r: TTGAAATTATAACTATAAA	f: TTTATAGTTGTAATTTCAA r: TTGAAATTACAACACTATAAA
Exon 5	AF136837.1	#42	f: CGAAGTTACATTAGGGAGA r: TCTCCCTAAATGTAACITCG	f: CGAAGTTACGTTAGGGAGA r: TCTCCCTAACTGTAACITCG
Exon 5	AF136837.1	#101	f: TCGGGACTTTATCGATTGC r: GCAATCGATAAAGTCCCGA	f: TCGGGACTTGATCGATTGC r: GCAATCGATCAAGTCCCG ₂
Exon 5	AF136837.1	#104	f: GGACTTTATCGATTGCTTC r: GAAAGCAATCGATAAAGTCC	f: GGACTTTATGGATTGCTTC r: GAAAGCAATCCATAAAGTCC
Exon 5	AF136837.1	#117	f: TGCTTCCTGATCAAAAATGG r: CCATTTTGAATCAGGAAGCA	f: TGCTTCCTGTTCAAAAATGG r: CCATTTTGAACAGGAAGCA
Exon 6	AF136838.1	#309	f: CACTTCTAGGAAAAGGACA r: TGTCCCTTTCTCTAGTTGTG	f: CACTTCTAGTAAAAGGACA r: TGTCCCTTTTACTAGTTGTG
Exon 7	NM_000770.1	#1135	f: GTCCCCCACC ₁ GGTGTGCCCC r: GGGGCACACCGGTGGGGAC	f: GTCCCCCACCAGTGTGCCCC r: GGGGCACACCTGGTGGGGAC
Exon 7	AF136840.1	#232	f: AGGATAGGAGCCACATGCC	f: AGGATAGGATCCACATGCC

Exon 8	AF136842.1	#206	r:GGCATGTGGCCTCCTATCCT	r:GGCATGTGGA/CTCCTATCCT	f:GGCATGTGGA A TCCTATCCT
			f:ATGATGACAAAGAATTTC	f:ATGATGACAA/GAGAAATTTC	f:ATGATGACAGAGAATTTC
Exon 8	AF136843.1	#30	r:GGAAATCTTTGTGCATCAT	r:GGAAATCTC/TTGTGCATCAT	r:GGAAATCTCTGTGCATCAT
			f:TGACCCCTGGCCACTTTCTA	f:TGACCCCTGGC/TCACTTTCTA	f:TGACCCCTGGTCACTTTCTA
Exon 8	AF136843.1	#87	r:TAGAAAGTGCCAGGGTCA	r:TAGAAAGTGA/GCCAGGGTCA	r:TAGAAAGTGA A CCAGGGTCA
			f:GCCCTTTCTCAAGCAGGTAAT	f:GCCCTTTCTCA/GGCAGGTAAT	f:GCCCTTTCTC G GCAGGTAAT
Intron 8	AF136843.1	#167	r:ATTACCTGCTGAGAAAGGC	r:ATTACCTGCC/TGAGAAAGGC	r:ATTACCTGCC C GGAGAAAGGC
			f:TACATGGCACTCCTCTGG	f:TACATGGCAC/ACTCCTCTGG	f:TACATGGCA A CTCCTCTGG
Intron 8	AF136843.1	#197	r:CCAGAGGAGTGCCATGTA	r:CCAGAGGAGT/GTGCCATGTA	r:CCAGAGGAGT T TGCCATGTA
			f:TTGCTATTGTCTCCATGATC	f:TTGCTATTTG/ATCCATGATC	f:TTGCTATTT A TCCATGATC
Intron 8	AF136843.1	#212	r:GATCATGGA C AAATAGCAA	r:GATCATGGAT/CAAATAGCAA	r:GATCATGGAT T AAATAGCAA
			f:GATCAAGAGCACCCTCTT	f:GATCAAGAGC/TACCCTCTT	f:GATCAAGAGT T ACCCTCTT
Intron 8	AF136843.1	#221	r:AAGAGTGGTGCTCTTGATC	r:AAGAGTGGTA/GCTCTTGATC	r:AAGAGTGGT A CTCTTGATC
			f:CACCACTCTTAACACCCAT	f:CACCACTCTT/CAACACCCAT	f:CACCACTCT C CAACACCCAT
Intron 8	AF136843.1	#255	r:ATGGGTGTTAAGAGTGGTG	r:ATGGGTGTG/AAGAGTGGTG	r:ATGGGTGT T GAGAGTGGTG
			f:AATACACCA T CAATTATTGG	f:AATACACCA T /CCATTATTGG	f:AATACACCA C CAATTATTGG
Intron 8	AF136843.1	#271	r:CCAATAATGATGGTGTATT	r:CCAATAATGG/ATGGTGTATT	r:CCAATAATG T TGGTGTATT
			f:TGGGCCAGATAGCGGGGCT	f:TGGGCCAGAT/CAGCGGGGCT	f:TGGGCCAGAC A AGCGGGGCT
Intron 8	AF136843.1	#271	r:AGCCCCGCTATCTGGCCCA	r:AGCCCCGCTG/ATCTGGCCCA	r:AGCCCCGCT T CTGGCCCA

Intron 8	AF136844.1	#118	f: TTATTTTACTGCATATTCTG r: CAGAAATATGCAGTAAATAA	f: TTATTTTACTACATATTCTG r: CAGAAATATGTAGTAAATAA
3' UTR	AF136845.1	#44	f: TCTGGCTGCCGATCTGCTA r: TAGCAGATCGCAGCCAGA	f: TCTGGCTGCTGATCTGCTA r: TAGCAGATCAGCAGCCAGA

Table 3 Comparison of allelic frequencies (%) from the populations analyzed (calculation based on Hardy-Weinberg law).

SNP (- : rel. to ATG) (GenBank Acc. No ref. to, s. text)	Caucasian	Japanese	African-American	Location
(- 1731) - (- 1732)	1,3	n.d.	11,7	5'UTR
-1627	n.d.	n.d.	16,6	5'UTR
-1478	n.d.	n.d.	2	5'UTR
-1325	n.d.	n.d.	1,4	5'UTR
-1221	n.d.	1,6	n.d.	5'UTR
-1214	4,3	n.d.	n.d.	5'UTR
-1207	13,5	n.d.	3,75	5'UTR
-1159	n.d.	n.d.	1,1	5'UTR
-1152	3,05	n.d.	n.d.	5'UTR
- 980	n.d.	n.d.	1,2	5'UTR
(- 765) - (- 767)	n.d.	n.d.	n.d.	5'UTR
(- 640) - (- 641)	10,35	n.d.	4,5	5'UTR
- 411	14	37,8	17,4	5'UTR
- 370	19,2	29,7	3,2	5'UTR
- 271	23,3	5,7	3,2	5'UTR
- 248	n.d.	n.d.	3,2	5'UTR
- 151	n.d.	n.d.	1,6	5'UTR
- 133	n.d.	n.d.	2,2	5'UTR
- 86	n.d.	n.d.	1,2	5'UTR
171	1,4	n.d.	n.d.	Intron 1
258	n.d.	n.d.	1,6	Intron 1
122*				Intron 2
150	n.d.	n.d.	11,6	Intron 2
180 -181	30,8	51,1	47,8	Intron 2
182	n.d.	n.d.	2,3	Intron 2
270	12	n.d.	4,5	Exon 3
334	n.d.	n.d.	15,9	Exon 3
378	14,6	n.d.	4,5	Intron 3
87	6,1	1,3	3,2	Intron 3
162	1,2	n.d.	n.d.	Intron 3
163	24,2	2,6	25,8	Intron 3
243 [§]				Exon 4
13	n.d.	n.d.	1,6	Exon 4
180	24,4	1,2	26,4	Intron 4
116	29,6	46,4	7,4	Intron 4
132	4,7	n.d.	7,4	Intron 4
172	24,6	51,2	24,1	Intron 4

189	2,6	n.d.	n.d.	Intron 4
42	n.d.	n.d.	1,9	Exon 5
101	n.d.	1,1	n.d.	Exon 5
104	5	n.d.	1,7	Exon 5
117	n.d.	n.d.	14,2	Exon 5
1135	n.d.	n.d.	1,2	Exon 7
206	11,4	n.d.	3,2	Exon 8
30	n.d.	7,3	n.d.	Exon 8
87	n.d.	n.d.	2,17	Exon 8
167	6,3	n.d.	5,6	Intron 8
197	23	51,2	41,1	Intron 8
212	1,3	n.d.	n.d.	Intron 8
221	n.d.	3,7	n.d.	Intron 8
255	n.d.	n.d.	1,1	Intron 8
271	2,1	n.d.	1,7	Intron 8
118	26,4	n.d.	44,4	Intron 8
44	22,3	53	n.d.	3'UTR

* - has been detected in a sample of a pre-screen (Caucasian sample).

§ - has been detected in a phenotyped sample (Caucasian sample).

n.d.- not detect in the samples analyzed.

Table 4 Listing of all amino acid changes in the coding regions

Met Gly Lys Arg Ser Ile Glu	s001.txt	Exon 3
Met Gly Lys Lys Ser Ile Glu	s002.txt	
Glu Leu Arg Lys Thr Lys Ala	s376.txt	Exon 3/4
Glu Leu Arg Lys Pro Arg Leu	s377.txt	
Arg Lys Thr Lys Ala Ser Pro	s003.txt	Exon 3/4
Arg Lys Thr Lys Ala Ser Pro	s004.txt	
Ile Cys Ser Val Val Phe Gln	s005.txt	Exon 3
Ile Cys Ser Ile Val Phe Gln	s006.txt	
Ala Ile Leu Asn Ser Pro Trp	s007.txt	Exon 4
Ala Ile Leu Ser Ser Pro Trp	s008.txt	
Arg Ser Tyr Ile Arg Glu Lys	s009.txt	Exon 5
Arg Ser Tyr Val Arg Glu Lys	s010.txt	
Pro Arg Asp Phe Ile Asp Cys	s011.txt	Exon 5
Pro Arg Asp Leu Ile Asp Cys	s012.txt	
Arg Asp Phe Ile Asp Cys Phe	s013.txt	Exon 5
Arg Asp Phe Met Asp Cys Phe	s014.txt	
Cys Phe Leu Ile Lys Met Glu	s015.txt	Exon 5
Cys Phe Leu Phe Lys Met Glu	s016.txt	
Met Glu Gln Glu Lys Asp Asn	s378.txt	Exon 6
Met Glu Gln STOP	s379.txt	
Val Pro Thr Gly Val Pro His	s017.txt	Exon 7
Val Pro Thr Ser Val Pro His	s018.txt	
Gln Asp Arg Ser His Met Pro	s380.txt	Exon 7
Gln Asp Arg Ile His Met Pro	s381.txt	
His Asp Asp Lys Glu Phe Pro	s019.txt	Exon 8
His Asp Asp Arg Glu Phe Pro	s020.txt	
Phe Asp Pro Gly His Phe Leu	s021.txt	Exon 8
Phe Asp Pro Gly His Phe Leu	s022.txt	
Met Pro Phe Ser Ala Gly Lys	s023.txt	Exon 8
Met Pro Phe Ser Ala Gly Lys	s024.txt	

CLAIMS

1. A polynucleotide comprising a polynucleotide selected from the group consisting of:
 - (a) a polynucleotide having the nucleic acid sequence of SEQ ID NO: 54, 57, 60, 63, 66, 69, 72, 75, 78, 81, 84, 87, 90, 93, 96, 99, 102, 105, 108, 111, 114, 117, 120, 123, 126, 129, 132, 135, 138, 141, 144, 147, 150, 153, 156, 159, 162, 165, 168, 171, 174, 177, 180, 183, 183, 189, 192, 195, 198, 201, 210, 213, 216, 219, 222, 225, 228, 231, 234, 237, 240, 243, 246, 249, 252, 255, 258, 261, 264, 267, 270, 273, 276, 279, 282, 285, 288, 291, 306, 309, 318, 321, 324, 327, 330, 333, 342, 345, 348, 351, 354, 357, 360, 363, 366, 369, 384, 387, 390, 393, 396 or 399;
 - (b) a polynucleotide encoding a polypeptide having the amino acid sequence of SEQ ID NO: 6, 8, 10, 12, 18, 377, 379 or 381;
 - (c) a polynucleotide capable of hybridizing to a CYP2C8 gene, wherein said polynucleotide is having at a position corresponding to position 411, 560, 713, 817, 824, 831, 879, 886, 1058, 1627, 1668, 1767, 1887, 1905 or 1952 (GenBank accession No: AF136830.1), at a position corresponding to position 171 or 258 (GenBank accession No: AF136832.1), at a position corresponding to position 122, 150, 182, 334, 339 or 378 (GenBank accession No: AF136833.1), at a position corresponding to position 162, 163, 243 (GenBank accession No: AF136834.2) or at position 583 (GenBank accession No: NM_000770.1), at a position corresponding to position 13 or 180 (GenBank accession No: AF136835.1), at a position corresponding to position 116, 132, 172 or 189 (GenBank accession No: AF136836.1), at a position corresponding to position 42 or 101 (GenBank accession No: AF136837.1), at a position corresponding to position 309 (GenBank accession No: AF136838.1), at a position corresponding to position 1135 (GenBank accession No: NM_000770.1), at a position corresponding to position 232 (GenBank accession No: AF136840.1), at a position corresponding to position 206 (GenBank accession No: AF136842.1), at a position corresponding to position 30, 87, 167, 197, 212, 221, 255 or 271 (GenBank accession No: AF136843.1), at a

position corresponding to position 118 (GenBank accession No: AF136844.1), at a position corresponding to position 44 (GenBank accession No: AF136845.1) of the cytochrome 2C8 gene (GenBank accession No: GI: 13787189) a nucleotide substitution, at a position corresponding to position 306 to 307, 1271 to 1273 or 1397 to 1398 of the CYP2C8 gene (GenBank accession No: AF136830.1), at a position corresponding to position 329 of the CYP2C8 gene (GenBank accession No: AF136833.1), at a position corresponding to position 87 of the CYP2C8 gene (GenBank accession No: AF136834.2) a deletion of one or more nucleotides or at a position corresponding to position 1785/1786 of the CYP2C8 gene (GenBank accession No: AF136830.1) or at a position corresponding to position 180/181 of the CYP2C8 gene (GenBank accession No: AF136833.1) an insertion of one or more nucleotides;

- (d) a polynucleotide capable of hybridizing to a CYP2C8 gene, wherein said polynucleotide is having at a position corresponding to position 411, 817, 824, 831, 879, 1058, 1767 or 1887 of the CYP2C8 gene (GenBank accession No: AF136830.1) an A, at a position corresponding to position 560 or 1668 of the CYP2C8 gene (GenBank accession No: AF136830.1) a G, at a position corresponding to position 713 or 886 of the CYP2C8 gene (GenBank accession No: AF136830.1) a T, at a position corresponding to position 1627, 1905 or 1952 of the CYP2C8 gene (GenBank accession No: AF136830.1) a C, at a position corresponding to position 258 of the CYP2C8 gene (GenBank accession No: AF136832.1) a T, at a position corresponding to position 171 of the CYP2C8 gene (GenBank accession No: AF136832.1) a C, at a position corresponding to position 122, 150 or 334 of the CYP2C8 gene (GenBank accession No: AF136833.1) an A, at a position corresponding to position 182 or 378 of the CYP2C8 gene (GenBank accession No: AF136833.1) a C, at a position corresponding to position 162, 163, 243 [identical to position corresponding to position 583 of the CYP2C8 gene (GenBank accession No: NM_000770.1) of the CYP2C8 gene (GenBank accession No: AF136834.2) an A, at a position

corresponding to position 180 of the CYP2C8 gene (GenBank accession No: AF136835.1) an A, at a position corresponding to position 13 of the CYP2C8 gene (GenBank accession No: AF136835.1) a G, at a position corresponding to position 116 or 132 of the CYP2C8 gene (GenBank accession No: AF136836.1) a G, at a position corresponding to position 172 of the CYP2C8 gene (GenBank accession No: AF136836.1) a G, at a position corresponding to position 189 of the CYP2C8 gene (GenBank accession No: AF136836.1) a C, at a position corresponding to position 42 or 101 of the CYP2C8 gene (GenBank accession No: AF136837.1) a G, at a position corresponding to position 1135 of the CYP2C8 gene (GenBank accession No: GI: 13787189) an A, at a position corresponding to position 309 of the CYP2C8 gene (GenBank accession No: AF136838.1) a T, at a position corresponding to position 232 (GenBank accession No: 136840.1) a T, at a position corresponding to position 30 or 212 of the CYP2C8 gene (GenBank accession No: AF136843.1) a T, at a position corresponding to position 87 of the CYP2C8 gene (GenBank accession No: AF136843.1) a G, at a position corresponding to position 167 or 197 of the CYP2C8 gene (GenBank accession No: AF136843.1) an A, at a position corresponding to position 221, 255 or 271 of the CYP2C8 gene (GenBank accession No: AF136843.1) a C, at a position corresponding to position 118 of the CYP2C8 gene (GenBank accession No: AF136844.1) an A, at a position corresponding to position 44 of the CYP2C8 gene (GenBank accession No: AF136845.1) a T;

- (e) a polynucleotide encoding a molecular CYP2C8 variant polypeptide or fragment thereof, wherein said polypeptide comprises an amino acid substitution at a position corresponding to any one of position 159, 181, 209, 244, 263, 274, 343 or 365 of the CYP2C8 polypeptide (GI: 13787189); and
- (f) a polynucleotide encoding a molecular CYP2C8 variant polypeptide or fragment thereof, wherein said polypeptide comprises an amino acid substitution of T to P at position corresponding to position 159 (frameshift), V to I at a position corresponding to position 181, N to S at

a position corresponding to position 209, I to V at a position corresponding to position 244, F to L at a position corresponding to position 263, E to Stop at a position corresponding to position 274, G to S at a position corresponding to position 365 or S to I at a position corresponding to position 343 of the CYP2C8 polypeptide (GenBank accession No: GI: 13787189).

2. A polynucleotide of claim 1, wherein said polynucleotide is associated with an incompatibility or disease related to arachidonic acid metabolism, cancer, cardiovascular diseases.
3. A polynucleotide of any one of claims 1 or 2 which is DNA or RNA.
4. A gene comprising the polynucleotide of any one of claims 1 or 2.
5. The gene of claim 4, wherein a nucleotide deletion, addition and/or substitution results in altered expression of the variant gene compared to the corresponding wild type gene.
6. A vector comprising a polynucleotide of any one of claims 1 to 3 or the gene of claim 4 or 5.
7. The vector of claim 6, wherein the polynucleotide is operatively linked to expression control sequences allowing expression in prokaryotic or eukaryotic cells or isolated fractions thereof.
8. A host cell genetically engineered with the polynucleotide of any one of claims 1 to 3, the gene of claim 4 or 5 or the vector of claim 6 or 7.
9. A method for producing a molecular variant CYP2C8 polypeptide or fragment thereof comprising
 - (a) culturing the host cell of claim 8; and
 - (b) recovering said protein or fragment from the culture.

10. A method for producing cells capable of expressing a molecular variant CYP2C8 polypeptide comprising genetically engineering cells with the polynucleotide of any one of claims 1 to 3, the gene of claim 4 or 5 or the vector of claim 6 or 7.
11. A polypeptide or fragment thereof encoded by the polynucleotide of any one of claims 1 to 3, the gene of claim 4 or 5 or obtainable by the method of claim 9 or from cells produced by the method of claim 10.
12. An antibody which binds specifically to the polypeptide of claim 11.
13. The antibody of claim 12 which specifically recognizes an epitope containing one or more amino acid substitution(s) resulting from a nucleotide exchange as defined in claim 1 or 5.
14. The antibody of claim 12 or 13 which is monoclonal or polyclonal.
15. A transgenic non-human animal comprising at least one polynucleotide of any one of claims 1 to 3, the gene of claim 4 or 5 or the vector of claim 6 or 7.
16. The transgenic non-human animal of claim 15 which is a mouse, a rat or a zebrafish.
17. A solid support comprising one or a plurality of the polynucleotide of any one of claims 1 to 3, the gene of claim 4 or 5, the vector of claim 6 or 7, the polypeptide of claim 11, the antibody of claim 12 or 13 or the host cell of claim 8 in immobilized form.
18. The solid support of claim 17, wherein said solid support is a membrane, a glass-or polypropylene- or silicon-chip, are oligonucleotide-conjugated beads or a bead array, which is assembled on an optical filter substrate.

19. An in vitro method for identifying a single nucleotide polymorphism said method comprising the steps of:
- (a) isolating a polynucleotide of any one claims 1 to 3 or the gene of claim 4 or 5 from a plurality of subgroups of individuals, wherein one subgroup has no prevalence for a CYP2C8 associated disease and at least one or more further subgroup(s) do have prevalence for a CYP2C8 associated disease; and
 - (b) identifying a single nucleotide polymorphism by comparing the nucleic acid sequence of said polynucleotide or said gene of said one subgroup having no prevalence for a CYP2C8 associated disease with said at least one or more further subgroup(s) having a prevalence for a CYP2C8 associated disease.
20. A method for identifying and obtaining a pro-drug or a drug capable of modulating the activity of a molecular variant of a CYP2C8 polypeptide comprising the steps of:
- (a) contacting the polypeptide of claim 11, the solid support of claim 17 or 18, a cell expressing a molecular variant gene comprising a polynucleotide of any one of claims 1 to 3, the gene of claim 4 or 5 or the vector of claim 6 or 7 in the presence of components capable of providing a detectable signal in response to drug activity with a compound to be screened for pro-drug or drug activity; and
 - (b) detecting the presence or absence of a signal or increase or decrease of a signal generated from the pro-drug or the drug activity, wherein the absence, presence, increase or decrease of the signal is indicative for a putative pro-drug or drug.
21. A method for identifying and obtaining an inhibitor of the activity of a molecular variant of a CYP2C8 polypeptide comprising the steps of:
- (a) contacting the protein of claim 11, the solid support of claim 17 or 18 or a cell expressing a molecular variant gene comprising a polynucleotide of any one of claims 1 to 3 or the gene of claim 4 or 5 or the vector of claim 6 or 7 in the presence of components capable of providing a

- detectable signal in response to drug activity with a compound to be screened for inhibiting activity; and
- (b) detecting the presence or absence of a signal or increase or decrease of a signal generated from the inhibiting activity, wherein the absence or decrease of the signal is indicative for a putative inhibitor.
22. The method of claim 20 or 21, wherein said cell is a cell of claim 8, obtained by the method of claim 10 or can be obtained by the transgenic non-human animal of claim 15 or 16.
23. A method of identifying and obtaining a pro-drug or drug capable of modulating the activity of a molecular variant of a CYP2C8 polypeptide comprising the steps of:
- (a) contacting the host cell of claim 8, the cell obtained by the method of claim 10, the polypeptide of claim 11 or the solid support of claim 17 or 18 with the first molecule known to be bound by a CYP2C8 polypeptide to form a first complex of said polypeptide and said first molecule;
- (b) contacting said first complex with a compound to be screened, and
- (c) measuring whether said compound displaces said first molecule from said first complex.
24. A method of identifying and obtaining an inhibitor capable of modulating the activity of a molecular variant of a CYP2C8 polypeptide or its gene product comprising the steps of:
- (a) contacting the host cell of claim 8, the cell obtained by the method of claim 10, the protein of claim 11 or the solid support of claim 17 or 18 with the first molecule known to be bound by a CYP2C8 polypeptide to form a first complex of said polypeptide and said first molecule;
- (b) contacting said first complex with a compound to be screened, and
- (c) measuring whether said compound displaces said first molecule from said first complex.

25. The method of claim 23 or 24, wherein said measuring step comprises measuring the formation of a second complex of said polypeptide and said compound.
26. The method of any one of claims 23 to 25, wherein said measuring step comprises measuring the amount of said first molecule that is not bound to said polypeptide.
27. The method of any one of claims 23 to 26, wherein said first molecule is labeled.
28. A method for the production of a pharmaceutical composition comprising the steps of the method of any one of claims 20 to 27; and the further step of formulating the compound identified and obtained or a derivative thereof in a pharmaceutically acceptable form.
29. A method of diagnosing a disorder related to the presence of a molecular variant of a CYP2C8 gene or susceptibility to such a disorder comprising determining the presence of a polynucleotide of any one of claims 1 to 3 or the gene of claim 4 or 5 in a sample from a subject.
30. The method of claim 29 further comprising determining the presence of a polypeptide of claim 11 or the antibody of any one of claims 12 to 14.
31. A method of diagnosing a disorder related to the presence of a molecular variant of a CYP2C8 gene or susceptibility to such a disorder comprising determining the presence of a polypeptide of claim 11 or the antibody of any one of claims 12 to 14 in a sample from a subject.
32. The method of any one of claims 29 to 31, wherein said disorder is a incompatibility or a disease related to arachidonic acid metabolism, cancer or cardiovascular diseases.

33. The method of any one of claims 29 to 32 comprising PCR, ligase chain reaction, restriction digestion, direct sequencing, nucleic acid amplification techniques, hybridization techniques or immunoassays.
34. A method of detection of the polynucleotide of any one of claims 1 to 3 or the gene of claim 4 or 5 in a sample comprising the steps of
 - (a) contacting the solid support of claim 17 or 18 with the sample under conditions allowing interaction of the polynucleotide of claim 1 to 3 or the gene of claim 4 or 5 with the immobilized targets on a solid support and
 - (b) determining the binding of said polynucleotide or said gene to said immobilized targets on a solid support.
35. An in vitro method for diagnosing a disease comprising the steps of the method of claim 34, wherein binding of said polynucleotide or gene to said immobilized targets on said solid support is indicative for the presence or the absence of said disease or a prevalence for said disease.
36. A diagnostic composition comprising the polynucleotide of any one of claims 1 to 3, the gene of claim 4 to 5, the vector of claim 6 or 7, the polypeptide of claim 11 or the antibody of any one of the claims 12 to 14.
37. A pharmaceutical composition comprising the polynucleotide of any one of claims 1 to 3, the gene of claim 4 or 5, the vector of claim 6 or 7, the polypeptide of claim 11 or the antibody of any of the claims 12 to 14.
38. Use the polynucleotide of any one of claims 1 to 3, the gene of claim 4 or 5, the vector of claim 6 or 7, the polypeptide of claim 11, or the antibody of any of the claims 12 to 14 for the preparation of a diagnostic composition for diagnosing a disease.
39. Use the polynucleotide of any one of claims 1 to 3, the gene of claim 4 or 5, the vector of claim 6 or 7, the polypeptide of claim 11, or the antibody of any of

the claims 12 to 14 for the preparation of a pharmaceutical composition for treating a disease.

40. The use of claims 38 or 39 , wherein said disease is an incompatibility or disease related to arachidonic acid metabolism, cancer or cardiovascular diseases.
41. A diagnostic kit for detection of a single nucleotide polymorphism comprising the polynucleotide of any one of claims 1 to 3, the gene of claim 4 or 5, the vector of claim 6 or 7, the polypeptide of claim 11, the antibody of any of the claims 12 to 14, the host cell of claim 8, the transgenic non-human animal of claim 15 or 16 or the solid support of claim 17 or 18.






Figure 1**Figure 2****Figure 3**

Figure 4**Figure 5**






Fig. 6

Fig. 7

AF136830.1 GI:6707879

```

cattttaatc cactgggtgct agaattatta actaaattaa tgtttatattt gaaagtcact 61
gattagatta atccacaagt attgaatttt agtcaatctt ggtggcccggt ttttaactgga 121
tgttttgctt aaaaggaagg cagcaagatg caggggttat ggtttccagc cccagcttgg 181
tcacttgcat tctgtgtgtc cttagctaaa gtactgaatc tccatggtct aactttctcc 241
tctctaaact gggaataaatt ttacagtggg caaagataat tgagagaata aaaagagatg 301
tgatgagtgt gaaaattctc tgtaaatttg tcataatgtc tataaacata atcgataaaa 361
cattgtataa ctgggtctaa tattttctta atgaaagagc tggaaataac tgtactggtc 421
aatttagaat aaaggtaatc ttttcagagc atgcctttgt atacacactt tgttattagt 481
gatctagttaa tggtcataaa tccagtgtga ttttagatctt catgaccatt gactatcagt 541
tcccatttca ggtctgcaca ttgcagtggg tctgtgccct ggggccattc agtgatttcc 601
ctgtgttcca tcttctgttg aatccacaac tgttgttctg tgtataattt ctcttcttg 661
ctgtgtatga ttacattcta ttatttgtaa caataacaga ccaaaaacaa tagaagcagc 721
catgtctgga ggtgactgga aggtggagaa gccatagatt ttcaagccct gtgccataaa 781
ttatgtgaga ttggcccttt ccttaatagt gctgaacaac tttcacttgt gaggtgatgc 841
agaggggaga actctaattt ttatttcttc ttttgagcgt ctccgggtct cttatcctta 901
taaacaataa acggacttct atttaatgtg aagcctgttg ctttctgaac agagtcaagg 961
tggcgtatct tcagagtaac taatgtctgg gggtgtttt gtttttctaa aattgttctt 1021
gagccagctg tggtgtaagt ggtaatgaac cccaatgggt atcagaagat ctctgctcaa 1081
atcccggttt taccggcaat gagctgtgtg gcactgacag gtgtcctggt ctcccagagt 1141
ttctttccca atttgaaaaa taaaaaatga taatctttat actccagtct cttttaatga 1201
tgaatataca ttatatata tacttttata tatttaatat aatatttaat agtataaata 1261
tgtatttatg ttattattat gtaataatgt atgtaacact ccctgctaata taacacaaat 1381
tctttgacat gtaaagtaaa taatcaccta ttattataat aatgtaataa taacacaaat 1381
attattatgt aataacatat atatttatgt atattgttta tatacattta aatatatata 1441
aatatacat tattagctaa taatttgata tatgtatggt aattcaacat gtatgagtta 1501
tattcactat ttcagtttta ggcagctgta ttttaagtga actatactaa atatttgaaa 1561
ggcttttgtt atcaagggtc aagtctccta ttttttgata tagcattaca atgtacattt 1621
tttatacaca aaatatagaa tacactgatt tccctcaagg tcataaattc ccaactgggtc 1681
attaatctga gaatattgaa ttttgagtat attctaacat agaatcattt acttcagtgt 1741
ttctccatca tcacagcaca ttggaacaac cagggacttt taattaaaaa tacctggggt 1801
ccaatccaat acaattaaac cagaatctcc tagattggca ctggaaagaa ggagtaggac 1861
aaaagaacat tttatttcta tccatggggt aaagtcact cagaaaaaaa gtataaattg 1921
gatctagggt attgtttact ttacatgtca aagagacaca cactaaatta gcagggagtg 1981
ttataaaaaa tttggagtg cagctcacag ctgtcttaat aagaagagaa ggtttcaatg 2041
gaaccttttg tggtcctggg gctgtgtctc tcttttatgc ttctcttttc actctgg

```

AF136831.1 GI:6707880

```

cagagctgta ggagaaggaa gctccctcct ggccccactc ctcttcctat tattggaaat 61
atgctacaga tagatgttaa ggacatctgc aaatctttca ccaatgtaag tctgccttat 121
gttccctccag ccaattgcaa agggtaagtt atttgactgc tattttttaga caaaatataat 181
tcctggaagc acactattat aatagatcat tgtaaagcaa aatactccct ctgaacttct 241
ttgatgtttc ttttgtcttc ctattttttt ttttttttga gacggagtct cgctctattg 301
cccaggctgg agtgacgtgg cactatctcc actcactgca agctctgcct cccaggttca 361
caccattctc ctgcctcagc ctcccccgag taactgggac tacaggtgcc ctccaccatg 421
cccggtcaat tttttgtatg

```

AF136832.1 GI:6707881

```

agtttcttca tttttaaaaa aggtcaaattg aatgtgctga atgtgttgaa gtgaggatga 61
actgtgtgat ttgtgtacca attgcctggg tcattgcgtg gcacatcaca ggccatctat 121
aagtggcagc tataacaatc accatcacat ttatgtacaa aattcagaaa tatcgaatct 181
atgtgtggca aatatgaaca ttaaaaaata caatgaaaat gtcagtctga atcatacata 241
gtatttggag caaatagcga cttatttttg tgctattttg atttcctttc ccagtttctca 301
aaagtctatg gtccgtgtgt caccgtgtat tttggcatga atcccatagt ggtgtttcat 361
ggatatgagg cagtgaagga agc

```

AF136833.1 GI:6707882

tactaaagga	cttggtaggt	gcacatat	tttggtaact	ggggtgaggg	61
ggatggaaaa	cagagcccta	aaaagcttct	cagcagagct	tagcctatct	121
cgagtgttgc	agcactttct	tccttggctg	tgaattctcc	cagtttctgc	181
attaggaatc	atttccagca	atggaaagag	atgggaaggag	atccggcggt	241
aaccttgcgg	aatttttgga	tggggaagag	gagcattgag	gaccgtgttc	301
tcactgcctt	gtggaggagt	tgagaaaaac	caagggtggg	tgactctact	361
gaccttaaca	gttacctgtc	ttcactagt	acgtccttgg	aaacatttca	421
ggtcttcatt	gogcatcctg	gttgtcagcc	ctcaggtggt	gga	

AF136834.2 GI:7544370

atgctcaact	catattttaag	gtaaaagtaa	tgtgttttatt	tcattccatgc	61
tggacacatg	gggaatttgt	aagatatgtt	taaaatttct	aaatttccct	121
cagatgcaaa	tctttttaaat	atttattttt	taataatttt	tttaaaaatt	181
tagcttcacc	ctgtgatccc	actttcatcc	tgggctgtgc	tccctgcaat	241
cag					

AF136835.1 GI:6707884

tcaggattct	gaactcccca	tggatccagg	taaggccaag	attttatttt	61
catttattca	agggtgtagg	gaagacttgg	tttaaaaatg	agaaaattga	121
cttttataca	ataaaaatga	tgtatgagtg	aagaaaataa	ttaccacctt	181
ttcaaaat	tcagcctcca	atcttttaggt	acagaaaatt	gctatatgtg	241
atttcccat	cagaagtgc	aggggtcagg	gaattccctt	tcctagccaa	301
gaacagatgg	cacctggaaa	attgggtcac	tcaccacctt	atactgtgct	361
tcttagtaaa					

AF136836.1 GI:6707885

tgggagatat	acctaattgta	tatgacgagt	tattgggtgc	agtacaccaa	61
gtatacatat	gtgacaaacc	tgcactttgt	gcacatatatac	cctagaactt	121
aaaaaatgta	tatatgtata	aaaatttccc	ttcaaaatgg	acatgatgtc	181
ttatagttat	aattttcaatc	agggcttggg	gtaagatata	tatatcttat	241
tatttaatat	tcttttctct	tttaggtctg	caataatttc	cctctactca	ttga

AF136837.1 GI:6707886

acaacaaagt	gcttaaaaat	gttgctctta	cacgaagtta	cattagggag	61
aacaccaagc	atcactggat	gttaacaatc	ctcgggactt	tatcgattgc	121
aatggagca	ggtaagatat	tagcaacaga	tcagtatttt	gatttcttgt	181
attcatcgaa	tccttctgt	atttactaag	gatgttttaa	tgatcaggcc	241
gacaagcatc	tttaattactt	attgtattta	tgggcctgca	ctaaacatca	301
aaaattgtcc	aatggctaga	atgcata			

AF136838.1 GI:6707887

ataacacaaa	ttgaagtaag	acagggcac	ggtatacttc	tgcttttatt	61
gaaatattct	gtgtgactaa	cctaagcagc	gaatgatttc	atgaatggaa	121
tgtcaggaaa	taaagtttga	gtcaactgat	ctgcagtttc	tgccatacca	181
ttttctaata	ctgtactgtc	cagtatctct	tttggctaac	tttaaaaaat	241
ttaaaattta	gtgtatttag	atatactggc	acataatttg	tcagataatt	301
acttctagga	aaaggacaac	caaaagtcag	aattcaatat	tgaaaa	

AF136839.1 GI:6707888

tcctgctcct	gctgaagcac	ccagaggtca	caggtaggac	cacagatgat	61
aatctcagaa	caatgctgag	aagatgggtc	cagtatcctc	caccttggtt	121
aggctcattc	tttaaatctt	tgtgtcatca	gctgtaatct	gtctaaattt	181
tttaaaatga	catctttgt				

AF136840.1 GI:6707889

tatattatgg	taattctttt	tatatggctg	gttggtacttc	tggacatgta	actcatgttt	61
gtaatgttgc	tgggattttt	atatcatgtt	aatgtggcca	tgaattgcta	tgacaaatgt	121
tccatatatc	ttcgtttcca	tcagttcttt	cttgtgtctt	gtcagctaaa	gtccaggaag	181
agattgatca	tgtaattggc	agacacagga	gcccctgcat	gcaggatagg	agccacatgc	241
cttacac						

AF136841.1 GI:6707890

gccccatgca	gtgaccactg	atactaagtt	cagaaactac	ctcatcccca	aggtaagctt	61
gtttctctta	cactatattt	ctgtacttct	gaaatttcca	tagtgctggg	ttgggttccaa	121
ccctctaaca	acacaagatg	agagaagtgc	aaaactcata	catgtggcag	cttga	

AF136842.1 GI:6707891

ccaccactgg	ccttaagctg	atccatgtaa	attactgtgt	ctggctggac	ctgagtttcc	61
tcattctatag	atcaacgtta	tggcgctacg	tgatgtccac	tacttctcct	cacttctgga	121
cttctttata	aatcagatta	tctgttttgt	tacttccagg	gcacaaccat	aatggcatta	181
ctgacttccg	tgctacatga	tgacaaagaa	tttcctaata	caaata		

AF136843.1 GI:6707892

tttcctaata	caaatatctt	tgaccctggc	cactttctag	ataagaatgg	caactttaag	61
aaaagtgact	acttcatgcc	tttctcagca	ggtaatagaa	actcgtttcc	atgtgtattt	121
aaaggaaaga	gagaactttt	tgggaattag	tgggaattac	atggcacctc	ctctggggct	181
ggtagaattg	ctatttgtcc	atgatcaaga	gcaccactct	taacacccat	gtgctccacc	241
ctcacaaata	accatcatta	ttggggccaga	tagcggggct	tgcaggagtt	aactctgttg	

AF136844.1 GI:6707893

aatatgtctc	ttttgtaca	tttgtttgtc	ccaccatcca	ttaatcaatc	catcatgtca	61
tccatccatt	catccacatg	ttcattcatt	tacccaatca	ttaatcaatt	atctactgca	121
tattctgttt	gtgcaagtca	caaataactg	tttgtcacag	tcacagttaa	acacaaggag	181
taactacttc	ctttctttgt	tatcttcagg	aaaacgaatt	tgtgcaggag	aaggacttgc	241
cgcgatgg						

AF136845.1 GI:6707894

tctgcttcat	ccctgtctga	agaatgctag	cccactctggc	tgccgatctg	ctatcacctg	61
caactctttt	tttatcaagg	acattcccac	tattatgtct	tctctgacct	ctcatcaaat	121
cttcccattc	actcaatata	ccataagcat	ccaaactcca	ttaaggagag	ttgttcagggt	181
cactgcacaa	atatactctg	aattattcat	actctgtaac	acttgattta	attgctgcat	241
atgctaatac	ttttctaata	ctgacttttt	aatatgttat	cactgtaaaa	cacagaaaag	

NM_000770.1 GI:4503220 mRNA

agtgcgaagt	cacagctgtc	ttaataagaa	gagaaggctt	caatggaacc	ttttgtgggtc	61
ctgggtgctgt	gtctctcttt	tatgcttctc	ttttcactct	ggagacagag	ctgtaggaga	121
aggaagctcc	ctcctggccc	cactcctctt	cctattattg	gaaatatgct	acagatagat	181
gttaaggaca	tctgcaaatc	tttcaccaat	ttctcaaaag	tctatgggtc	tgtgttcacc	241
gtgtattttg	gcatgaatcc	catagtgggtg	tttcatggat	atgaggcagt	gaagggaagcc	301
ctgattgata	atggagagga	gttttctgga	agaggcaatt	ccccaatata	tcaaagaatt	361
actaaaggac	ttggaatcat	ttccagcaat	ggaaagagat	ggaaggagat	ccggcggttcc	421
tccttcacaa	ccttgcgga	ttttgggatg	gggaagagga	gcattgagga	ccgtgttcaa	481
gaggaagctc	actgccttgt	ggaggagtgt	agaaaaacca	aggcttcacc	ctgtgatccc	541
actttcatcc	tgggctgtgc	tccttgcaat	gtgatctgct	ccgttggttt	ccagaaacga	601
tttgattata	aagatcagaa	ttttctcacc	ctgatgaaaa	gattcaatga	aaacttcagg	661
attctgaact	ccccatggat	ccagggtctgc	aataatttcc	ctctactcat	tgattgtttc	721
ccaggaactc	acaacaaagt	gcttaaaaat	gttgctctta	cacgaagtta	cattagggag	781
aaagtaaaag	aacaccaaag	atcactggat	gttaacaatc	ctcgggactt	tatogattgc	841
ttcctgatca	aaatggagca	ggaaaaggac	aaccaaaagt	cagaattcaa	tattgaaac	901
ttgggttgga	ctgtagctga	tctatttgtt	gctggaacag	agacaacaag	caccactctg	961
agatatggac	tcctgctcct	gctgaagcac	ccagagggtca	cagctaaagt	ccaggaagag	1021
attgatcatg	taattggcag	acacaggagc	ccctgcatgc	aggataggag	ccacatgcct	1081
tacactgatg	ctgtagtgc	cgagatccag	agatacagtg	accttggtcc	cacgggtgtg	1141
ccccatgcag	tgaccactga	tactaagttc	agaaaactacc	tcaccccaa	gggcacaacc	1201
ataatggcat	tactgacttc	cgtgctacat	gatgacaaag	aatttcctaa	tccaaatata	1261
tttgaccctg	gccactttct	agataagaat	ggcaacttta	agaaaagtga	ctacttcatg	1321


```
cctttctcag caggaaaacg aattttgtgca ggagaaggac ttgcccgcac ggagctatatt 1381
ttattttctaa ccacaatttt acagaacttt aacctgaaat ctgttgatga tttaaagaac 1441
ctcaataacta ctgcagttac caaagggatt gtttctctgc caccctcata ccagatctgc 1501
ttcatccctg tctgaagaat gctagcccat ctggctgctg atctgctatc acctgcaact 1561
cttttttttat caaggacatt cccactatta tgtcttctct gacctctcat caaatcttcc 1621
cattcactca atatcccata agcatccaaa ctccattaag gagagtgtgt caggtcactg 1681
cacaaatata tctgcaatta ttcatactct gtaacacttg tattaattgc tgcataatgct 1741
aatactttttc taatgctgac tttttaatat gttatcactg taaaacacag aaaagtgatt 1801
aatgaatgat aatttagatc catcttcttt gtgaatgtgc taaataaaaa gtgttattaa 1861
ttgcta
```

NP_000761.2 GI:13787189

```
MEPFVVLVLC LSFMLLFSLW RQSCRRRKLP PGPTPLPIIG NMLQIDVKDI CKSFTNFSKV 61
YGPVFTVYFG MNPIVVFHGY EAVKEALIDN GEEFSGRGNS PISQRITKGL GISSNGKRW 121
KEIRRFSLTN LRNFGMGKRS IEDRVQEEAH CLVEELRKTG ASPCDPTFIL GCAPCNVICS 181
VVFQKRFDYK DQNFLTLMKR FNENFRILNS PWIQVCNNFP LLIDCFPGTH NKVLKNVALT 241
RSYIREKVKE HQASLDVNNP RDFMDCFLIK MEQEKDNQKS EFNIENTLVGT VADLFVAGTE 301
TTSTTLRYGL LLLLKHPEVT AKVQEEIDHV IGRHRSPCMQ DRSHPYTDV VVHEIQRYSD 361
LVPTGVPHAV TTDTKFRNYL IPKGTTIMAL LTSVLHDDKE FPNPNIFDPG HFLDKNGNFK 421
KSDYFMPFSA GKRICAGEGL ARMELFLFLT TILQNFNLKS VDDLKNLNTT AVTKGIVSLP 481
PSYQICFIPV
```


SEQUENCE LISTING

<110> EPIDAUROS Biotechnologie AG
PENGGER, Anja
SPRENGER, Reimund
BRINKMANN, Ulrich

<120> Polymorphisms in the human gene for cytochrome P450
polypeptide 2C8 and their use in diagnostic and
therapeutic applications

<130> F 1754 PCT

<150> EP 01 11 2899.8

<151> 2001-06-01

<160> 417

<170> PatentIn Ver. 2.1

<210> 1

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 1

Met Gly Lys Arg Ser Ile Glu
1 5

<210> 2

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 2

Met Gly Lys Lys Ser Ile Glu
1 5

<210> 3

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 3

Arg Lys Thr Lys Ala Ser Pro
1 5

<210> 4
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 4
Arg Lys Thr Lys Ala Ser Pro
1 5

<210> 5
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 5
Ile Cys Ser Val Val Phe Gln
1 5

<210> 6
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 6
Ile Cys Ser Ile Val Phe Gln
1 5

<210> 7
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 7
Ala Ile Leu Asn Ser Pro Trp
1 5

<210> 8
<211> 7
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 8

Ala Ile Leu Ser Ser Pro Trp
1 5

<210> 9

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 9

Arg Ser Tyr Ile Arg Glu Lys
1 5

<210> 10

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 10

Arg Ser Tyr Val Arg Glu Lys
1 5

<210> 11

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 11

Pro Arg Asp Phe Ile Asp Cys
1 5

<210> 12

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 12

Pro Arg Asp Leu Ile Asp Cys
1 5

<210> 13

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 13

Arg Asp Phe Ile Asp Cys Phe
1 5

<210> 14

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 14

Arg Asp Phe Met Asp Cys Phe
1 5

<210> 15

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 15

Cys Phe Leu Ile Lys Met Glu
1 5

<210> 16

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 16

Cys Phe Leu Phe Lys Met Glu
1 5

<210> 17
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 17
Val Pro Thr Gly Val Pro His
1 5

<210> 18
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 18
Val Pro Thr Ser Val Pro His
1 5

<210> 19
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 19
His Asp Asp Lys Glu Phe Pro
1 5

<210> 20
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 20
His Asp Asp Arg Glu Phe Pro
1 5

<210> 21
<211> 7
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 21

Phe Asp Pro Gly His Phe Leu
1 5

<210> 22

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 22

Phe Asp Pro Gly His Phe Leu
1 5

<210> 23

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 23

Met Pro Phe Ser Ala Gly Lys
1 5

<210> 24

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 24

Met Pro Phe Ser Ala Gly Lys
1 5

<210> 25

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 25
atcttagtca atcttggtgg cccg 24

<210> 26
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 26
ttcaacagaa gatggaacac aggga 25

<210> 27
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 27
tcattgacct tgactatcag ttccc 25

<210> 28
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 28
tgatacccat tggggttcat tacc 24

<210> 29
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 29
aacagagtca aggtggcgta tcttc 25

<210> 30
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 30
caatattctc agattaatga ccagttggg 29

<210> 31
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 31
agacttagcc ctgataaca aaagcc 26

<210> 32
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 32
gttttaggcag ctgtatttta agtgaac 27

<210> 33
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 33
actccaaagt tttataaca ctccc 25

<210> 34
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 34
ggcactggaa agaaggagta ggac 24

<210> 35

<211> 27
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 35
gatctattat aatagtgtgc ttccagg

27

<210> 36
<211> 22
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 36
ttgtgtacca attgcctggg tc

22

<210> 37
<211> 24
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 37
tttttagggc tctgttttcc atcc

24

<210> 38
<211> 24
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 38
gagcttagcc tatctgcatg gctg

24

<210> 39
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 39

acctggccac ccctgaaatg

20

<210> 40

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 40

tccatgctga ttttttttgg acac

24

<210> 41

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 41

ctgacccctt gcacttctga tg

22

<210> 42

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 42

tgacgagtta ttgggtgcag tacac

25

<210> 43

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 43

ttccatgatg tttagtgacg gcc

23

<210> 44

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 44

ttgaagtaag acagggcatc gg

22

<210> 45

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 45

agaaacaagg tggaggatac tggc

24

<210> 46

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 46

ggccatgaat tgctatgaca aatg

24

<210> 47

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 47

ggttggaacc aaaccagcac tatg

24

<210> 48

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 48

ctggctggac ctgagtttcc tc

22

<210> 49

<211> 20

<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 49

ttaactcctg caagccccgc

20

<210> 50

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 50

gtacatttgt ttgtcccacc atcc

24

<210> 51

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 51

tgcagtgacc tgaacaactc tcct

24

<210> 52

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 52

gatgtgatga gtgtgaaaat

20

<210> 53

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> n=ag or deleted

<400> 53
gatgtgatgn tgtgaaaat 19

<210> 54
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 54
gatgtgatgt gtgaaaat 18

<210> 55
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 55
atatttcacac tcatcacatc 20

<210> 56
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> n=ct or deleted

<400> 56
atatttcacan catcacatc 19

<210> 57
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 57
atatttcaca catcacatc 18

<210> 58

<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 58
ggaaataact gtactggtc

19

<210> 59
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> w=a or t

<400> 59
ggaaataacw gtactggtc

19

<210> 60
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 60
ggaaataaca gtactggtc

19

<210> 61
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 61
gaccagtaca gttatttcc

19

<210> 62
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial

sequence

<220>

<223> w=a or t

<400> 62

gaccagtacw gttattttcc

19

<210> 63

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 63

gaccagtact gttattttcc

19

<210> 64

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 64

ggtctgcaca ttgcagtgg

19

<210> 65

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> r=g or a

<400> 65

ggtctgcacr ttgcagtgg

19

<210> 66

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 66

ggtctgcacg ttgcagtgg

19

<210> 67

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 67

ccactgcaat gtgcagacc

19

<210> 68

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> y=t or c

<400> 68

ccactgcaay gtgcagacc

19

<210> 69

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 69

ccactgcaac gtgcagacc

19

<210> 70

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 70

aaaacaatag aagcagcca

19

<210> 71

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> k=g or t

<400> 71

aaaacaatak aagcagcca

19

<210> 72

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 72

aaaacaatat aagcagcca

19

<210> 73

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 73

tggctgcttc tattgtttt

19

<210> 74

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> m=a or c

<400> 74

tggctgcttm tattgtttt

19

<210> 75

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 75

tggctgctta tattgtttt

19

<210> 76

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 76

agtgtgaac aactttcac

19

<210> 77

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> m=a or c

<400> 77

agtgtgaam aactttcac

19

<210> 78

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 78

agtgtgaaa aactttcac

19

<210> 79

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 79

gtgaaagttg ttcagcact

19

<210> 80
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> k=g or t

<400> 80
gtgaaagttk ttcagcact

19

<210> 81
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 81
gtgaaagttt ttcagcact

19

<210> 82
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 82
aacaactttc acttgtgag

19

<210> 83
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> m=a or c

<400> 83
aacaactttm acttgtgag

19

<210> 84
<211> 19

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 84
aacaacttta acttgtag

19

<210> 85
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 85
ctcacaagtg aaagttggt

19

<210> 86
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> k=g or t

<400> 86
ctcacaagtk aaagttggt

19

<210> 87
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 87
ctcacaagtt aaagttggt

19

<210> 88
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 88
ttcacttggtg aggtgatgc

19

<210> 89
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> r=g or a

<400> 89
ttcacttgtr aggtgatgc

19

<210> 90
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 90
ttcacttgta aggtgatgc

19

<210> 91
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 91
gcatcacctc acaagtga

19

<210> 92
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> y=t or c

<400> 92
gcatcaccty acaagtga

19

<210> 93
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 93
gcatcacctt acaagtga

19

<210> 94
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 94
cttttgagcg tctccggtc

19

<210> 95
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> r=g or a

<400> 95
cttttgagcr tctccggtc

19

<210> 96
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 96
cttttgagca tctccggtc

19

<210> 97
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 97

gaccggagac gctcaaaag

19

<210> 98

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> y=t or c

<400> 98

gaccggagay gctcaaaag

19

<210> 99

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 99

gaccggagat gctcaaaag

19

<210> 100

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 100

gcgtctccgg tcctcttat

19

<210> 101

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> k=g or t

<400> 101

gcgtctccgk tcctcttat

19

<210> 102

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 102

gcgtctccgt tcctcttat

19

<210> 103

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 103

ataagaggac cggagacgc

19

<210> 104

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> m=a or c

<400> 104

ataagaggam cggagacgc

19

<210> 105

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 105

ataagaggaa cggagacgc

19

<210> 106
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 106
accccaatgg gtatcagaa

19

<210> 107
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> r=g or a

<400> 107
accccaatgr gtatcagaa

19

<210> 108
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 108
accccaatga gtatcagaa

19

<210> 109
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 109
ttctgatacc cattggggt

19

<210> 110
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> y=t or c

<400> 110

ttctgatacy cattggggt

19

<210> 111

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 111

ttctgatact cattggggt

19

<210> 112

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 112

gtatttatgt tattattatg t

21

<210> 113

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> n=tta or deleted

<400> 113

gtatttatgn ttattatgt

19

<210> 114

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 114
gtattttatgt tattatgt

18

<210> 115
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 115
acataataat aacataaata c

21

<210> 116
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> n=taa or deleted

<400> 116
acataataan cataaatac

19

<210> 117
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 117
acataataac ataaatac

18

<210> 118
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 118
tgtaataaca tatatatatta

20

<210> 119

<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> n=at or deleted

<400> 119
tgtaataacn atatatatta

19

<210> 120
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 120
tgtaataaca tatatttta

18

<210> 121
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 121
taaatatata tgttattaca

20

<210> 122
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> n=at or deleted

<400> 122
taaatatatn gttattaca

19

<210> 123
<211> 18
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 123

taaatatatg ttattaca

18

<210> 124

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 124

ttttttatat acaaaatat

19

<210> 125

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> y=t or c

<400> 125

ttttttatay acaaaatat

19

<210> 126

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 126

ttttttatac acaaaatat

19

<210> 127

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 127

atatttttgta tataaaaaaa

19

<210> 128

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> r=g or a

<400> 128

atattttgtr tataaaaaaa

19

<210> 129

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 129

atattttgtg tataaaaaaa

19

<210> 130

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 130

ggtcataaat tcccaactg

19

<210> 131

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> k=g or t

<400> 131

ggtcataaak tcccaactg

19

<210> 132
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 132
ggtcataaag tcccaactg

19

<210> 133
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 133
cagttgggaa tttatgacc

19

<210> 134
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> m=a or c

<400> 134
cagttgggam tttatgacc

19

<210> 135
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 135
cagttgggac tttatgacc

19

<210> 136
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 136

acattggaac aaccaggga

19

<210> 137

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220> cora

<223> y=t or c

<400> 137

acattggaay aaccaggga

19

<210> 138

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 138

acattggaaa aaccaggga

19

<210> 139

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 139

tccttggttg ttccaatgt

19

<210> 140

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220> gart

<223> r=g or a

<400> 140
tccctgggttr ttccaatgt 19

<210> 141
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 141
tccctgggttt ttccaatgt 19

<210> 142
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 142
aattaaaaat acctgggc 18

<210> 143
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> n=a or aa

<400> 143
aattaaaant acctgggc 18

<210> 144
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 144
aattaaaaaa tacctgggc 19

<210> 145
<211> 18

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 145
gccaggtat ttttaatt

18

<210> 146
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> n=a or at

<400> 146
gccaggtnt ttttaatt

18

<210> 147
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 147
gccaggtat tttttaatt

19

<210> 148
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 148
ctatccatgg gccaaagtc

19

<210> 149
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> r=g or a

<400> 149

ctatccatgr gccaaagtc

19

<210> 150

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 150

ctatccatga gccaaagtc

19

<210> 151

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 151

gactttggcc catggatag

19

<210> 152

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> y=t or c

<400> 152

gactttggcy catggatag

19

<210> 153

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 153

gactttggct catggatag

19

<210> 154
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 154
ccactcagaa aaaaagtat

19

<210> 155
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> m=a or c

<400> 155
ccactcagam aaaaagtat

19

<210> 156
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 156
ccactcagac aaaaagtat

19

<210> 157
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 157
atactttttt tctgagtgg

19

<210> 158
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> k=g or t

<400> 158

atactttttk tctgagtgg

19

<210> 159

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 159

atactttttg tctgagtgg

19

<210> 160

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 160

acatgtcaaa gagacacac

19

<210> 161

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> m=a or c

<400> 161

acatgtcaam gagacacac

19

<210> 162

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial

sequence

<400> 162
acatgtcaac gagacacac 19

<210> 163
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 163
gtgtgtctct ttgacatgt 19

<210> 164
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> k=g or t

<400> 164
gtgtgtctck ttgacatgt 19

<210> 165
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 165
gtgtgtctcgt ttgacatgt 19

<210> 166
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 166
attcagaaat atcgaatct 19

<210> 167
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> y=t or c

<400> 167
attcagaaay atcgaatct

19

<210> 168
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 168
attcagaaac atcgaatct

19

<210> 169
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 169
agattcgata tttctgaat

19

<210> 170
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> r=g or a

<400> 170
agattcgatr tttctgaat

19

<210> 171
<211> 19
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 171

agattcgatg tttctgaat

19

<210> 172

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 172

agcaaatagc gacttattt

19

<210> 173

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> y=t or c

<400> 173

agcaaatagy gacttattt

19

<210> 174

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 174

agcaaatagt gacttattt

19

<210> 175

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 175
aaataagtcg ctatttgct

19

<210> 176
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> r=g or a

<400> 176
aaataagtcg ctatttgct

19

<210> 177
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 177
aaataagtca ctatttgct

19

<210> 178
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 178
atggctgccg agtggttgca

19

<210> 179
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> r=g or a

<400> 179
atggctgccc agtggttgca

19

<210> 180
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 180
atggctgcca agtgttgca

19

<210> 181
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 181
tgcaacactc ggcagccat

19

<210> 182
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> y=t or c

<400> 182
tgcaacacty ggcagccat

19

<210> 183
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 183
tgcaacactt ggcagccat

19

<210> 184
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 184

tccttggctg tgaattctc

19

<210> 185

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> r=g or a

<400> 185

tccttggctr tgaattctc

19

<210> 186

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 186

tccttggcta tgaattctc

19

<210> 187

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 187

gagaattcac agccaagga

19

<210> 188

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> y=t or c

<400> 188
gagaattcay agccaagga 19

<210> 189
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 189
gagaattcat agccaagga 19

<210> 190
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 190
ccttttttta ttaggaat 18

<210> 191
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> n=t or tt

<400> 191
ccttttttna ttaggaat 18

<210> 192
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 192
cctttttttt attaggaat 19

<210> 193

<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 193
attcctaata aaaaaagg

18

<210> 194
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> n=a or aa

<400> 194
attcctaata aaaaaagg

18

<210> 195
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 195
attcctaata aaaaaagg

19

<210> 196
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 196
cttttttttat taggaatca

19

<210> 197
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial

sequence

<220>

<223> y=t or c

<400> 197

ctttttttay taggaatca

19

<210> 198

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 198

ctttttttac taggaatca

19

<210> 199

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 199

tgattcctaa taaaaaaag

19

<210> 200

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> r=g or a

<400> 200

tgattcctar taaaaaaag

19

<210> 201

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 201

tgattcctag taaaaaaag

19

<210> 202

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 202

tggggaagag gagcattga

19

<210> 203

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> r=g or a

<400> 203

tggggaagar gagcattga

19

<210> 204

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 204

tggggaagaa gagcattga

19

<210> 205

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 205

tcaatgctcc tcttcccca

19

<210> 206

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> y=t or c

<400> 206

tcaatgctcy tcttcccca

19

<210> 207

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 207

tcaatgctct tcttcccca

19

<210> 208

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 208

aaaaaccaag ggtgggtga

19

<210> 209

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> r=g or a

<400> 209

aaaaaccaar ggtgggtga

19

<210> 210

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 210

aaaaaccaa ggtgggtga

19

<210> 211

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 211

tcacccaccc ttggttttt

19

<210> 212

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> y=t or c

<400> 212

tcacccaccc ttggttttt

19

<210> 213

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 213

tcacccaccc ttggttttt

19

<210> 214

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 214

cagttacctg tcttcacta

19

<210> 215
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> s=g or c

<400> 215
cagttaccts tcttcacta

19

<210> 216
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 216
cagttacctc tcttcacta

19

<210> 217
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 217
tagtgaagac aggtaactg

19

<210> 218
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> s=g or c

<400> 218
tagtgaagas aggtaactg

19

<210> 219
<211> 19

<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 219
tagtgaagag aggtaactg

19

<210> 220
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 220
tgtaagatat gtttaaaat

19

<210> 221
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> n=t or deleted

<400> 221
tgtaagatan gtttaaaat

19

<210> 222
<211> 18
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 222
tgtaagatag tttaaaat

18

<210> 223
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 223
atatttaaaca tatctttaca

19

<210> 224
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> n=a or deleted

<400> 224
atatttaaact tatctttaca

19

<210> 225
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 225
atatttaaact atctttaca

18

<210> 226
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 226
ataattttttt taaaaattt

19

<210> 227
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> w=a or t

<400> 227
ataattttttw taaaaattt

19

<210> 228
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 228
ataatttttta taaaaattt 19

<210> 229
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 229
aaattttttaa aaaaatttat 19

<210> 230
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> w=a or t

<400> 230
aaattttttaw aaaaatttat 19

<210> 231
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 231
aaattttttat aaaaatttat 19

<210> 232
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 232
taattttttt aaaaatttt

19

<210> 233
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> w=a or t

<400> 233
taattttttw aaaaatttt

19

<210> 234
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 234
taatttttta aaaaatttt

19

<210> 235
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 235
aaaattttta aaaaatta

19

<210> 236
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> w=a or t

<400> 236

aaaattttttw aaaaaatta

19

<210> 237

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 237

aaaattttttt aaaaaatta

19

<210> 238

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 238

atctgctccg ttgttttcc

19

<210> 239

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> r=g or a

<400> 239

atctgctccr ttgttttcc

19

<210> 240

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 240

atctgctcca ttgttttcc

19

<210> 241
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 241
ggaaaacaac ggagcagat

19

<210> 242
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> y=t or c

<400> 242
ggaaaacaay ggagcagat

19

<210> 243
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 243
ggaaaacaat ggagcagat

19

<210> 244
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 244
ggattctgaa ctccccatg

19

<210> 245
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> r=g or a

<400> 245

ggattctgar ctcccatg

19

<210> 246

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 246

ggattctgag ctcccatg

19

<210> 247

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 247

catggggagt tcagaatcc

19

<210> 248

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> y=t or c

<400> 248

catggggagy tcagaatcc

19

<210> 249

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 249
catgggggagc tcagaatcc 19

<210> 250
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 250
tgatttcctg ttcaaaatt 19

<210> 251
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> r=g or a

<400> 251
tgatttcctr ttcaaaatt 19

<210> 252
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 252
tgatttccta ttcaaaatt 19

<210> 253
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 253
aattttgaac aggaaatca 19

<210> 254
<211> 19

<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> y=t or c

<400> 254

aattttgaay aggaaatca

19

<210> 255

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 255

aattttgaat aggaaatca

19

<210> 256

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 256

acttaaagta taataaaaa

19

<210> 257

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> r=g or a

<400> 257

acttaaagtr taataaaaa

19

<210> 258

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 258

acttaaagtg taataaaaa

19

<210> 259

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 259

tttttattat actttaagt

19

<210> 260

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> y=t or c

<400> 260

tttttattay actttaagt

19

<210> 261

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 261

tttttattac actttaagt

19

<210> 262

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 262

aaaatgtata tatgtataa

19

<210> 263
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> r=g or a

<400> 263
aaaatgtatr tatgtataa

19

<210> 264
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 264
aaaatgtatg tatgtataa

19

<210> 265
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 265
ttatacatat atacatttt

19

<210> 266
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> y=t or c

<400> 266
ttatacatay atacatttt

19

<210> 267

<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 267
ttatacatatc atacatttt

19

<210> 268
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 268
atgatgtctt attcatatt

19

<210> 269
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> y=t or c

<400> 269
atgatgtcty attcatatt

19

<210> 270
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 270
atgatgtctc attcatatt

19

<210> 271
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial

sequence

<400> 271
aatatgaata agacatcat 19

<210> 272
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> r=g or a

<400> 272
aatatgaatr agacatcat 19

<210> 273
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 273
aatatgaatg agacatcat 19

<210> 274
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 274
tttatagtta taatttcaa 19

<210> 275
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> r=g or a

<400> 275

tttatagtttr taatttcaa

19

<210> 276

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 276

tttatagttg taatttcaa

19

<210> 277

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 277

ttgaaattat aactataaa

19

<210> 278

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> y=t or c

<400> 278

ttgaaattay aactataaa

19

<210> 279

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 279

ttgaaattac aactataaa

19

<210> 280

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 280

cgaagttaca ttagggaga

19

<210> 281

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> r=g or a

<400> 281

cgaagttacr ttagggaga

19

<210> 282

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 282

cgaagttacg ttagggaga

19

<210> 283

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 283

tctccctaata gtaacttcg

19

<210> 284

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> y=t or c

<400> 284

tctccctaay gtaacttcg

19

<210> 285

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 285

tctccctaac gtaacttcg

19

<210> 286

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 286

tcgggacttt atcgattgc

19

<210> 287

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> k=g or t

<400> 287

tcgggacttk atcgattgc

19

<210> 288

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 288

tcgggacttg atcgattgc

19

<210> 289
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 289
gcaatcgata aagtcccga

19

<210> 290
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> m=a or c

<400> 290
gcaatcgatm aagtcccga

19

<210> 291
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 291
gcaatcgatc aagtcccga

19

<210> 292
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 292
ggactttatc gattgcttc

19

<210> 293
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> s=g or c

<400> 293

ggactttats gattgcttc

19

<210> 294

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 294

ggactttatg gattgcttc

19

<210> 295

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 295

gaagcaatcg ataaagtcc

19

<210> 296

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> s=g or c

<400> 296

gaagcaatcs ataaagtcc

19

<210> 297

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 297
gaagcaatcc ataaagtcc 19

<210> 298
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 298
tgcttcctga tcaaaatgg 19

<210> 299
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> w=a or t

<400> 299
tgcttcctgw tcaaaatgg 19

<210> 300
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 300
tgcttcctgt tcaaaatgg 19

<210> 301
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 301
ccattttgat caggaagca 19

<210> 302

<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> w=a or t

<400> 302
ccattttgaw caggaagca

19

<210> 303
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 303
ccattttgaa caggaagca

19

<210> 304
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 304
gtccccaccg gtgtgcccc

19

<210> 305
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> r=g or a

<400> 305
gtccccaccr gtgtgcccc

19

<210> 306
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 306
gtccccacca gtgtgcccc

19

<210> 307
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 307
gggggcacacc ggtggggac

19

<210> 308
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> y=t or c

<400> 308
gggggcacacy ggtggggac

19

<210> 309
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 309
gggggcacact ggtggggac

19

<210> 310
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 310

atgatgacaa agaatttcc

19

<210> 311

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> r=g or a

<400> 311

atgatgacar agaatttcc

19

<210> 312

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 312

atgatgacag agaatttcc

19

<210> 313

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 313

ggaaattctt tgatcatcat

19

<210> 314

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> y=t or c

<400> 314

ggaaattcty tgatcatcat

19

<210> 315
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 315
ggaaattctc tgatcatcat 19

<210> 316
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 316
tgaccctggc cacttttcta 19

<210> 317
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> y=t or c

<400> 317
tgaccctggy cacttttcta 19

<210> 318
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 318
tgaccctggt cacttttcta 19

<210> 319
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 319

tagaaagtgg ccaggggtca

19

<210> 320

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> r=g or a

<400> 320

tagaaagtgr ccaggggtca

19

<210> 321

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 321

tagaaagtga ccaggggtca

19

<210> 322

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 322

gcctttctca gcaggtaat

19

<210> 323

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> r=g or a

<400> 323
gcctttctcr gcaggtaat

19

<210> 324
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 324
gcctttctcg gcaggtaat

19

<210> 325
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 325
attacctgct gagaaaggc

19

<210> 326
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> y=t or c

<400> 326
attacctgcy gagaaaggc

19

<210> 327
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 327
attacctgcc gagaaaggc

19

<210> 328
<211> 19

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 328
tacatggcac ctcctctgg

19

<210> 329
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> m=a or c

<400> 329
tacatggcam ctcctctgg

19

<210> 330
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 330
tacatggcaa ctcctctgg

19

<210> 331
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 331
ccagaggagg tgccatgta

19

<210> 332
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> k=g or t

<400> 332
ccagaggagk tgccatgta 19

<210> 333
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 333
ccagaggagt tgccatgta 19

<210> 334
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 334
ttgctatttg tccatgatc 19

<210> 335
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> r=g or a

<400> 335
ttgctatttr tccatgatc 19

<210> 336
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 336
ttgctattta tccatgatc 19

<210> 337
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 337
gatcatggac aaatagcaa

19

<210> 338
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> y=t or c

<400> 338
gatcatggay aaatagcaa

19

<210> 339
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 339
gatcatggat aaatagcaa

19

<210> 340
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 340
gatcaagagc accactctt

19

<210> 341
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> y=t or c

<400> 341

gatcaagagy accactcctt

19

<210> 342

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 342

gatcaagagt accactcctt

19

<210> 343

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 343

aagagtgggtg ctcttgatc

19

<210> 344

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<220>

<223> r=g or a

<400> 344

aagagtggtr ctcttgatc

19

<210> 345

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial

sequence

<400> 345
aagagtggta ctcttgatc 19

<210> 346
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 346
caccactctt aacacccat 19

<210> 347
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> y=t or c

<400> 347
caccactcty aacacccat 19

<210> 348
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 348
caccactctc aacacccat 19

<210> 349
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 349
atgggtgtta agagtgggtg 19

<210> 350
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> r=g or a

<400> 350
atgggtgttr agagtgggtg

19

<210> 351
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 351
atgggtgttg agagtgggtg

19

<210> 352
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 352
aatacaccat cattattgg

19

<210> 353
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> y=t or c

<400> 353
aatacaccay cattattgg

19

<210> 354
<211> 19
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 354

aatacaccac cattattgg

19

<210> 355

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 355

ccaataatga tgggtgtatt

19

<210> 356

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<220>

<223> r=g or a

<400> 356

ccaataatgr tgggtgtatt

19

<210> 357

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 357

ccaataatgg tgggtgtatt

19

<210> 358

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 358
tggggccagat agcggggct 19

<210> 359
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> y=t or c

<400> 359
tggggccagay agcggggct 19

<210> 360
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 360
tggggccagac agcggggct 19

<210> 361
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 361
agccccgcta tctggccca 19

<210> 362
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> r=g or a

<400> 362
agccccgctr tctggccca 19

<210> 363
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 363
agccccgctg tctggccca

19

<210> 364
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 364
ttattttactg catattctg

19

<210> 365
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> r=g or a

<400> 365
ttattttactr catattctg

19

<210> 366
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 366
ttattttacta catattctg

19

<210> 367
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 367
cagaatatgc agtaaataa 19

<210> 368
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> y=t or c

<400> 368
cagaatatgy agtaaataa 19

<210> 369
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 369
cagaatatgt agtaaataa 19

<210> 370
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 370
tctggctgcc gatctgcta 19

<210> 371
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> y=t or c

<400> 371
tctggctgcy gatctgcta 19

<210> 372
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 372
tctggctgct gatctgcta 19

<210> 373
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 373
tagcagatcg gcagccaga 19

<210> 374
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<220>
<223> r=g or a

<400> 374
tagcagatcr gcagccaga 19

<210> 375
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 375
tagcagatca gcagccaga 19

<210> 376

<211> 7
<212> PRT
<213> Homo sapiens

<400> 376
Glu Leu Arg Lys Thr Lys Ala
1 5

<210> 377
<211> 7
<212> PRT
<213> Homo sapiens

<400> 376
Glu Leu Arg Lys Pro Arg Leu
1 5

<210> 378
<211> 7
<212> PRT
<213> Homo sapiens

<400> 378
Met Glu Gln Glu Lys Asp Asn
1 5

<210> 379
<211> 3
<212> PRT
<213> Homo sapiens

<400> 379
Met Glu Gln
1

<210> 380
<211> 7
<212> PRT
<213> Homo sapiens

<400> 380
Gln Asp Arg Ser His Met Pro
1 5

<210> 381
<211> 7
<212> PRT
<213> Homo sapiens

<400> 381
Gln Asp Arg Ile His Met Pro
1 5

<210> 382
<211> 19

<212> DNA
<213> Homo sapiens

<400> 382
ttgagaaaaa ccaagggtg

19

<210> 383
<211> 18
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> 9...9
<223> n = a oder aa

<400> 383
ttgagaaanc caagggtg

18

<210> 384
<211> 18
<212> DNA
<213> Homo sapiens

<400> 384
ttgagaaaac caagggtg

18

<210> 385
<211> 19
<212> DNA
<213> Homo sapiens

<400> 385
cacccttggt ttttctcaa

19

<210> 386
<211> 18
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> 10...10
<223> n = t or tt

<400> 386
cacccttggn ttttctcaa

18

<210> 387
<211> 18
<212> DNA
<213> Homo sapiens

<400> 387
cacccttggt ttttctcaa

18

<210> 388
<211> 19
<212> DNA
<213> Homo sapiens

<400> 388
cacttctagg aaaaggaca

19

<210> 389
<211> 19
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> 10...10
<223> n = g or t

<400> 389
cacttctagn aaaaggaca

19

<210> 390
<211> 19
<212> DNA
<213> Homo sapiens

<400> 390
cacttctagt aaaaggaca

19

<210> 391
<211> 19
<212> DNA
<213> Homo sapiens

<400> 391
tgtccttttc ctagttgtg

19

<210> 392
<211> 19
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> 10...10
<223> n = c or a

<400> 392
tgtccttttn ctagttgtg

19

<210> 393
<211> 19
<212> DNA
<213> Homo sapiens

<400> 393
tgtcctttta ctagttgtg

19

<210> 394
<211> 19
<212> DNA
<213> Homo sapiens

<400> 394
aggataggag ccacatgcc

19

<210> 395
<211> 19
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> 10...10
<223> n = g or t

<400> 395
aggataggan ccacatgcc

19

<210> 396
<211> 19
<212> DNA
<213> Homo sapiens

<400> 396
aggataggat ccacatgcc

19

<210> 397
<211> 19
<212> DNA
<213> Homo sapiens

<400> 397
ggcatgtggc tcctatcct

19

<210> 398
<211> 19
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> 10...10
<223> n = a or c

<400> 398
ggcatgtggn tcctatcct

19

<210> 399
<211> 19

<212> DNA

<213> Homo sapiens

<400> 399

ggcatgtgga tcctatcct

19

<210> 400

<211> 2097

<212> DNA

<213> Homo sapiens

<220>

<223> genomic DNA

<400> 400

cattttaatc cactggtgct agaattatta actaaattaa tgtttatfff gaaagtcact	60
gattagatta atccacaagt attgaatfff agtcaatcct ggtggcccg tttactgga	120
tgttttgctt aaaaggaagg cagcaagatg caggggttat ggtttccagc cccagcttg	180
tcacttgcac tctgtgtgtc cttagctaaa gtactgaatc tccatggtct aactttctcc	240
tctctaaact gggaataatt ttacagtggg caaagataat tgagagaata aaaagagatg	300
tgatgagtgt gaaaattctc tgtaaatttg tcataatgtc tataaacata atcgataaaa	360
cattgtataa ctgggtctaa tattttctta atgaaagagc tggaaataac tgtactggtc	420
aatttagaat aaaggtaatc ttttcagagc atgcctttgt atacacactt tgttattagt	480
gatctagtaa tgttcataaa tccagttgta tttagatcct catgaccatt gactatcagt	540
tcccatttca ggtctgcaca ttgcagtggc tctgtgccct ggtccattc agtgatttcc	600
ctgtgttcca tcttctgttg aatccacaac tgttgttctg tgtataattt ctcttccttg	660
ctgtgtatga ttacattcta ttatttgtaa caataacaga ccaaaaacaa tagaagcagc	720
catgtctgga ggtgactgga aggtggagaa gccatagatt ttcaagccct gtgccataaa	780
ttatgtgaga ttggcccttt ccttaatagt gctgaacaac tttcacttgt gaggtgatgc	840
agagggggaga actctaattt ttatttcttc ttttgagcgt ctccggtcct cttatcctta	900
taaacaaata acggacttct atttaatgtg aagcctgttg ctttctgaac agagtcaagg	960
tggcgtatct tcagagtaac taatgtctgg ggtttgtttt gtttttctaa aattgttctt	1020
gagccagctg tgggtgtaagt ggtaatgaac cccaatgggt atcagaagat ctctgctcaa	1080
atccccggtt taccggcaat gagctgtgtg gcactgacag gtgtcctgtt ctcccagagt	1140
ttctttccca atttgaaaaa taaaaaatga taatctttat actccagtct cttttaatga	1200

tgaatataca	tttatatata	tactttttata	tattttaatat	aatattttaat	agtataaata	1260
tgtattttatg	ttattattat	gtaataatgt	atgtaacact	ccctgcta	tcagttttgtc	1320
tctttgacat	gtaaagtaaa	taatcaccta	ttattataat	aatgtaataa	taacacaaat	1380
attattatgt	aataacatat	atattttatgt	atattgttta	tatacattta	aatatatata	1440
aatatacatt	tattagctaa	taatttgata	tatgtatgg	aattcaacat	gtatgagtta	1500
tattcactat	ttcatgttta	ggcagctgta	ttttaagtga	actatactaa	atatttgaaa	1560
ggctttttgtt	atcaagggct	aagtctccta	ttttttgata	tagcattaca	atgtacattt	1620
tttatacaca	aaatatagaa	tacactgatt	tcctcaagg	tcataaattc	ccaactgggtc	1680
attaatctga	gaatattgaa	ttttgagtat	attctaacat	agaatcattt	acttcagtgt	1740
ttctccatca	tcacagcaca	ttggaacaac	cagggacttt	taattaaaaa	tacctgggct	1800
ccaatccaat	acaattaaac	cagaatctcc	tagattggca	ctggaaagaa	ggagtaggac	1860
aaaagaacat	tttattttcta	tccatgggcc	aaagtccact	cagaaaaaaa	gtataaattg	1920
gatctaggtg	attgtttact	ttacatgtca	aagagacaca	cactaaatta	gcagggagtg	1980
ttataaaaaac	tttggagtgc	aagctcacag	ctgtcttaat	aagaagagaa	ggcttcaatg	2040
gaaccttttg	tggtcctggt	gctgtgtctc	tcttttatgc	ttctcttttc	actctgg	2097

<210> 401

<211> 440

<212> DNA

<213> Homo sapiens

<220>

<223> genomic DNA

<400> 401

cagagctgta	ggagaaggaa	gctccctcct	ggccccactc	ctcttccctat	tattggaaat	60
atgctacaga	tagatgttaa	ggacatctgc	aaatctttca	ccaatgtaag	tctgccttat	120
gttccctccag	ccaattgcaa	agggttaagt	atttgactgc	tattttttaga	caaaatatat	180
tcctggaagc	acactattat	aatagatcat	tgtaaagcaa	aatactccct	ctgaacttct	240
ttgatgtttc	ttttgtcttc	ctattttttt	ttttttttga	gacggagtct	cgctctattg	300
cccaggctgg	agtgcagtgg	cactatctcc	actcactgca	agctctgcct	cccaggttca	360
caccattctc	ctgcctcagc	ctcccccgag	taactgggac	tacaggtgcc	ctccaccatg	420

cccggctaatt tttttgtatg

440

<210> 402

<211> 383

<212> DNA

<213> Homo sapiens

<220>

<223> genomic DNA

<400> 402

agttttcttca ttttttaaaac aggtcaaagt aatgtgctga atgtgttgaa gtgaggatga 60
actgtgtgat ttgtgtacca attgcctggg tcattgcgtg gcacatcaca ggccatctat 120
aagtggcagc tataacaatc accatcacat ttatgtacaa aattcagaaa tatcgaatct 180
atgtgtggca aatatgaaca ttaaaaaata caatgaaaat gtcagtctga atcatacata 240
gtatattggag caaatagcga cttatatttg tgctatattgc atttcctttc ccagttctca 300
aaagtctatg gtcctgtgtt caccgtgtat tttggcatga atcccatagt ggtgtttcat 360
ggatatgagg cagtgaagga agc 383

<210> 403

<211> 463

<212> DNA

<213> Homo sapiens

<220>

<223> genomic DNA

<400> 403

tactaaagga cttggtaggt gcacatattt ctgtgtcagc tttggttaact ggggtgaggg 60
ggatggaaaa cagagcccta aaaagcttct cagcagagct tagcctatct gcatggctgc 120
cgagtgttgc agcactttct tccttggctg tgaattctcc cagtttctgc cccttttttt 180
attaggaatc atttccagca atggaaagag atggaaggag atccggcggt tctccctcac 240
aaccttgcgg aattttggga tggggaagag gagcattgag gaccgtgttc aagaggaagc 300
tactgcctt gtggaggagt tgagaaaaac caaggtggg tgactctact ctgcgtcatt 360
gaccttaaca gttacctgtc ttcactagt acgtccttgg aaacatttca ggggtggcca 420

ggtcttcatt gcgcacctcg gttgtcagcc ctcagggtggt gga 463

<210> 404

<211> 243

<212> DNA

<213> Homo sapiens

<220>

<223> genomic DNA

<400> 404
atgctcaact catatttaag gtaaaagtaa tgtgtttatt tcatccatgc tgattttttt 60
tggacacatg gggaatttgt aagatatgtt taaaatttct aaatttcctt tatgtcttaa 120
cagatgcaaa tcttttaaatt atttattttt taataatttt tttaaaaatt tttaaattctt 180
tagcttcacc ctgtgatccc actttcatoc tgggctgtgc tccctgcaat gtgatctgct 240
ccg 243

<210> 405

<211> 370

<212> DNA

<213> Homo sapiens

<220>

<223> genomic DNA

<400> 405
tcaggattct gaactcccca tggatccagg taaggccaag attttatttt ccttggaac 60
cattttattca aggttgtagg gaagacttgg tttaaaaatg agaaaattga tactaaaatg 120
cttttataca ataaaaatga tgtatgagtg aagaaaataa ttaccacctt tgatttcctg 180
ttcaaaattt tcagcctcca atcttttaggt acagaaaatt gctatatgtg cacaataaaa 240
atttcccatc cagaagtgca aggggtcagg gaattccctt tcctagccaa gcaaagctgt 300
gaacagatgg cacctggaaa attgggtcac toccacccta atactgtgct tttctagtgg 360
tcttagtaaa 370

<210> 406

<211> 294

<212> DNA

<213> Homo sapiens

<220>

<223> genomic DNA

<400> 406

tgggagatat acctaattgta tatgacgagt tattgggtgc agtacaccaa cctggcacat	60
gtatacatat gtgacaaaacc tgcactttgt gcacatatat cctagaactt aaagtataat	120
aaaaaatgta tatatgtata aaaatttccc ttcaaatgg acatgatgtc ttattcatat	180
ttatagttat aatttcaatc agggcttgg gtaagatata tatatcttat gacatgttta	240
tatttaatat tcttttctct tttaggtctg caataatttc cctctactca ttga	294

<210> 407

<211> 327

<212> DNA

<213> Homo sapiens

<220>

<223> genomic DNA

<400> 407

acaacaaagt gcttaaaaat gttgctctta cacgaagtta cattagggag aaagtaaaag	60
aacaccaagc atcactggat gttaacaatc ctcgggactt tatcgattgc ttctgatca	120
aatggagca ggtaagatat tagcaacaga tcagtatttt gatctcttgc ccattttgtg	180
attcatcgaa tccttctgta atttactaag gatgttttaa tgatcaggcc agtaatgctt	240
gacaagcatc ttaattactt attgtattta tgggcctgca ctaaacaatca tggaaaatac	300
aaaattgtcc aatggctaga atgcata	327

<210> 408

<211> 346

<212> DNA

<213> Homo sapiens

<220>

<223> genomic DNA

<400> 408

ataacacaaa ttgaagtaag acagggcatc ggtatacttc tgctttttatt tctgggggaaa	60
gaaatattct gtgtgactaa cctaagcagc gaatgatttc atgaatggaa cttgtaggct	120
tgtcaggaaa taaagtttga gtcaactgat ctgcagtttc tgccatacca cacagttgct	180
ttttctaata ctgtactgtc cagtatctct tttggctaac tttaaaaaat agtatgtttt	240
ttaaaattta gtgtatttag atatactggc acataatttg tcagataatt gcatgaaatc	300
acttctagga aaaggacaac caaaagtcag aattcaatat tgaaaa	346

<210> 409

<211> 199

<212> DNA

<213> Homo sapiens

<220>

<223> genomic DNA

<400> 409

tcctgctcct gctgaagcac ccagaggcca caggtaggac cacagatgat gaacaaagtg	60
aatttcagaa caatgctgag aagatgggtc cagtatcctc caccttggtt ctctcagaga	120
aggctcattc tttaaatttc tgtgtcatca gctgtaatct gtctaaattt gatgacacaa	180
tttaaaatga catctttgt	199

<210> 410

<211> 247

<212> DNA

<213> Homo sapiens

<220>

<223> genomic DNA

<400> 410

tatattatgg taattctttt tatatggctg gttgtacttc tggacatgta actcatgttt	60
-------------------------------------------------------------------	----

gtaatgttgc tgggattttt atatcatggt aatgtggcca tgaattgcta tgacaaatgt 120
tccatatatc ttcgtttcca tcagttcttt cttgtgtctt gtcagctaaa gtccaggaag 180
agattgatca tgtaattggc agacacagga gcccctgcat gcaggatagg agccacatgc 240
cttacac 247

<210> 411

<211> 175

<212> DNA

<213> Homo sapiens

<220>

<223> genomic DNA

<400> 411

gccccatgca gtgaccactg atactaagtt cagaaactac ctcatcccca aggtaagctt 60
gtttctctta cactatattt ctgtacttct gaaatttcca tagtgctggg ttgggttccaa 120
ccctctaaca acacaagatg agagaagtgc aaaactcata catgtggcag cttega 175

<210> 412

<211> 226

<212> DNA

<213> Homo sapiens

<220>

<223> genomic DNA

<400> 412

ccaccactgg ccttaagctg atccatgtaa attactgtgt ctggctggac ctgagtttcc 60
tcattctatag atcaacgtta tggogctacg tgatgtccac tacttctcct cacttctgga 120
cttctttata aatcagatta tctgttttgt tacttccagg gcacaaccat aatggcatta 180
ctgacttccg tgctacatga tgacaaagaa tttcctaata caaata 226

<210> 413

<211> 300

<212> DNA

<213> Homo sapiens

<220>

<223> genomic DNA

<400> 413

tttcttaatc caaatatctt tgaccctggc cactttctag ataagaatgg caactttaag	60
aaaagtgact acttcatgcc tttctcagca ggtaatagaa actcgtttcc atttgtatct	120
aaaggaaaga gagaactttt tggaattagt tggaatttac atggcacctc ctctgggggt	180
ggtagaattg ctatttgtcc atgatcaaga gcaccactct taacacccat gtgctccacc	240
ctcacaatac accatcatta ttggggccaga tagcgggggct tgcaggagtt aactctgttg	300

<210> 414

<211> 248

<212> DNA

<213> Homo sapiens

<220>

<223> genomic DNA

<400> 414

aatatgtctc tttttgtaca ttgtttgtc ccaccatcca ttaatcaatc catcatgtca	60
tccatccatt catccacatg ttcatcctc taccatca ttaatcaatt atttactgca	120
tattctgttt gtgcaagtca caaatgactg ttgttcacag tcacagttaa acacaaggag	180
taactacttc ctttctttgt tatcttcagg aaaacgaatt tgtgcaggag aaggacttgc	240
ccgcatgg	248

<210> 415

<211> 300

<212> DNA

<213> Homo sapiens

<220>

<223> genomic DNA

<400> 415

tctgcttcat ccctgtctga agaatgctag cccatctggc tgccgatctg ctatcacctg 60
caactctttt tttatcaagg acattccac tattatgtct tctctgacct ctcatcaaatt 120
cttcccattc actcaatata ccataagcat ccaaactcca ttaaggagag ttgttcaggt 180
cactgcacaa atatatactgc aattattcat actctgtaac acttggtatta attgctgcat 240
atgctaatac ttttctaata ctgacttttt aatatgttat cactgtaaaa cacagaaaag 300

<210> 416

<211> 1866

<212> DNA

<213> Homo sapiens

<220>

<223> cDNA

<400> 416

agtgcgaagct cacagctgtc ttaataagaa gagaaggctt caatggaacc ttttgtggtc 60
ctgggtgctgt gtctctcttt tatgcttctc ttttactct ggagacagag ctgtaggaga 120
aggaagctcc ctcttgccc cactcctctt cctattattg gaaatatgct acagatagat 180
gttaaggaca tctgcaaata tttcaccaat ttctcaaaag tctatgggtc tgtgttcacc 240
gtgtattttg gcatgaatcc catagtgggtg tttcatggat atgaggcagt gaaggaagcc 300
ctgattgata atggagagga gttttctgga agaggcaatt cccaatata tcaaagaatt 360
actaaaggac ttggaatcat ttccagcaat ggaaagagat ggaaggagat ccggcgtttc 420
tccctcacia ccttgcgga ttttgggatg gggaagagga gcattgagga ccgtgttcaa 480
gaggaagctc actgccttgt ggaggagtgt agaaaaacca aggcttcacc ctgtgatccc 540
actttcatcc tgggctgtgc tccctgcaat gtgatctgct ccgttgtttt ccagaaacga 600
tttgattata aagatcagaa ttttctcacc ctgatgaaaa gattcaatga aaacttcagg 660
attctgaact cccatggat ccaggctctgc aataatttcc ctctactcat tgattgtttc 720
ccaggaactc acaacaaagt gcttaaaaat gttgctctta cacgaagtta cattagggag 780
aaagtaaaag aacaccaagc atcactggat gttaacaata ctcgggactt tatcgattgc 840
ttcctgatca aaatggagca ggaaaaggac aacaaaaagt cagaattcaa tattgaaaac 900
ttggttggca ctgtagctga tctatttggt gctggaacag agacaacaag caccactctg 960
agatatggac tctgctcct gctgaagcac ccagagggtc cagctaaagt ccaggaagag 1020
attgatcatg taattggcag acacaggagc cctgcatgc aggataggag ccacatgcct 1080


```

tacactgatg ctgtagtgca cgagatccag agatacagtg accttgtccc caccggtgtg 1140
cccatgcag tgaccactga tactaagttc agaaactacc tcatcccca gggcacaacc 1200
ataatggcat tactgacttc cgtgctacat gatgacaaag aatttcctaa tccaaatatc 1260
tttgaccctg gccactttct agataagaat ggcaacttta agaaaagtga ctacttcatg 1320
cctttctcag caggaaaacg aatttgtgca ggagaaggac ttgcccgcag ggagctatct 1380
ttattttctaa ccacaatttt acagaacttt aacctgaaat ctgttgatga tttaaagaac 1440
ctcaatacta ctgcagttac caaagggatt gtttctctgc caccctcata ccagatctgc 1500
ttcatccctg tctgaagaat gctagcccat ctggctgctg atctgctatc acctgcaact 1560
ctttttttat caaggacatt cccactatta tgtcttctct gacctctcat caaatcttcc 1620
cattcactca atatcccata agcatccaaa ctccattaag gagagttggt caggtcactg 1680
cacaaatata tctgcaatta ttcatactct gtaacacttg tattaattgc tgcataatgct 1740
aatacttttc taatgctgac tttttaatat gttatcactg taaaacacag aaaagtgatt 1800
aatgaatgat aatttagatc catttctttt gtgaatgtgc taaataaaaa gtgttattaa 1860
ttgcta 1866

```

<210> 417

<211> 490

<212> PRT

<213> Homo sapiens

<400> 417

```

Met Glu Pro Phe Val Val Leu Val Leu Cys Leu Ser Phe Met Leu Leu
1           5           10           15

```

```

Phe Ser Leu Trp Arg Gln Ser Cys Arg Arg Arg Lys Leu Pro Pro Gly
          20           25           30

```

```

Pro Thr Pro Leu Pro Ile Ile Gly Asn Met Leu Gln Ile Asp Val Lys
          35           40           45

```

```

Asp Ile Cys Lys Ser Phe Thr Asn Phe Ser Lys Val Tyr Gly Pro Val
          50           55           60

```

```

Phe Thr Val Tyr Phe Gly Met Asn Pro Ile Val Val Phe His Gly Tyr
65           70           75           80

```


Glu Ala Val Lys Glu Ala Leu Ile Asp Asn Gly Glu Glu Phe Ser Gly
 85 90 95

Arg Gly Asn Ser Pro Ile Ser Gln Arg Ile Thr Lys Gly Leu Gly Ile
 100 105 110

Ile Ser Ser Asn Gly Lys Arg Trp Lys Glu Ile Arg Arg Phe Ser Leu
 115 120 125

Thr Asn Leu Arg Asn Phe Gly Met Gly Lys Arg Ser Ile Glu Asp Arg
 130 135 140

Val Gln Glu Glu Ala His Cys Leu Val Glu Glu Leu Arg Lys Thr Lys
 145 150 155 160

Ala Ser Pro Cys Asp Pro Thr Phe Ile Leu Gly Cys Ala Pro Cys Asn
 165 170 175

Val Ile Cys Ser Val Val Phe Gln Lys Arg Phe Asp Tyr Lys Asp Gln
 180 185 190

Asn Phe Leu Thr Leu Met Lys Arg Phe Asn Glu Asn Phe Arg Ile Leu
 195 200 205

Asn Ser Pro Trp Ile Gln Val Cys Asn Asn Phe Pro Leu Leu Ile Asp
 210 215 220

Cys Phe Pro Gly Thr His Asn Lys Val Leu Lys Asn Val Ala Leu Thr
 225 230 235 240

Arg Ser Tyr Ile Arg Glu Lys Val Lys Glu His Gln Ala Ser Leu Asp
 245 250 255

Val Asn Asn Pro Arg Asp Phe Met Asp Cys Phe Leu Ile Lys Met Glu
 260 265 270

Gln Glu Lys Asp Asn Gln Lys Ser Glu Phe Asn Ile Glu Asn Leu Val
 275 280 285

Gly Thr Val Ala Asp Leu Phe Val Ala Gly Thr Glu Thr Thr Ser Thr
 290 295 300

Thr Leu Arg Tyr Gly Leu Leu Leu Leu Lys His Pro Glu Val Thr
 305 310 315 320

Ala Lys Val Gln Glu Glu Ile Asp His Val Ile Gly Arg His Arg Ser
 325 330 335

Pro Cys Met Gln Asp Arg Ser His Met Pro Tyr Thr Asp Ala Val Val
 340 345 350

His Glu Ile Gln Arg Tyr Ser Asp Leu Val Pro Thr Gly Val Pro His
 355 360 365

Ala Val Thr Thr Asp Thr Lys Phe Arg Asn Tyr Leu Ile Pro Lys Gly
 370 375 380

Thr Thr Ile Met Ala Leu Leu Thr Ser Val Leu His Asp Asp Lys Glu
 385 390 395 400

Phe Pro Asn Pro Asn Ile Phe Asp Pro Gly His Phe Leu Asp Lys Asn
 405 410 415

Gly Asn Phe Lys Lys Ser Asp Tyr Phe Met Pro Phe Ser Ala Gly Lys
 420 425 430

Arg Ile Cys Ala Gly Glu Gly Leu Ala Arg Met Glu Leu Phe Leu Phe
 435 440 445

Leu Thr Thr Ile Leu Gln Asn Phe Asn Leu Lys Ser Val Asp Asp Leu
 450 455 460

Lys Asn Leu Asn Thr Thr Ala Val Thr Lys Gly Ile Val Ser Leu Pro
 465 470 475 480

Pro Ser Tyr Gln Ile Cys Phe Ile Pro Val
 485 490